arXiv:1412.2924v1 [math.RT] 9 Dec 2014

UNITARY DUAL OF P-ADIC $U(5)$

CLAUDIA SCHOEMANN

Abstract. We study the parabolically induced complex representations of the unitary group in 5 variables, $U(5)$, defined over a p-adic field.

Let F be a p-adic field. Let $E : F$ be a field extension of degree two. Let $\mathrm{Gal}(E : F) = \{\mathrm{id}, \sigma\}$. We write $\sigma(x) = \overline{x}\ \forall x \in E$. Let $E^* := E \setminus \{0\}$ and let $E^1 := \{x \in E \mid x\overline{x} = 1\}$.

$U(5)$ has three proper standard Levi subgroups, the minimal Levi subgroup $M_0 \cong E^* \times E^* \times E^1$ and the two maximal Levi subgroups $M_1 \cong \mathrm{GL}(2, E) \times E^1$ and $M_2 \cong E^* \times U(3)$.

We consider representations induced from the minimal Levi subgroup M_0, representations induced from non-cuspidal, not fully-induced representations of the two maximal Levi subgroups M_1 and M_2, and representations induced from cuspidal representations of M_1.

We describe - except several particular cases - the unitary dual in terms of Langlands-quotients.

Contents

1. Introduction 1
2. Definitions 2
3. Irreducible unitary representations of $U(5)$, in terms of Langlands-quotients 4
3.1. Representations with cuspidal support in M_0, fully-induced 4
3.2. Representations induced from M_1, with cuspidal support in M_0, not fully-induced 17
3.3. Representations induced from M_2, with cuspidal support in M_0, not fully-induced 20
3.4. Representations with cuspidal support in M_1 23
References 28

1. Introduction

We study the parabolically induced complex representations of the unitary group in 5 variables - $U(5)$ - defined over a non-archimedean local field of characteristic 0. This is $\mathbb{Q}_p$ or a finite extension of $\mathbb{Q}_p$, where p is a prime number. We speak of a 'p-adic field'.

Let F be a p-adic field. Let $E : F$ be a field extension of degree two. Let $\mathrm{Gal}(E : F) = \{\mathrm{id}, \sigma\}$ be the Galois group. We write $\sigma(x) = \overline{x}\ \forall x \in E$. Let $E^* := E \setminus \{0\}$ and let $E^1 := \{x \in E \mid x\overline{x} = 1\}$. Let $|\ |_E$ denote the p-adic norm on E.

$U(5)$ has three proper parabolic subgroups. Let P_0 denote the minimal standard parabolic subgroup and P_1 and P_2 the two maximal standard parabolic subgroups. One has the Levi decomposition $P_i = M_i N_i,\ i \in \{0, 1, 2\}$, where M_i denote the standard Levi subgroups and N_i are the corresponding unipotent subgroups of $U(5)$.

Date: October 2014.

$M_0 \cong E^* \times E^* \times E^1$ is the minimal Levi subgroup, $M_1 \cong \mathrm{GL}(2,E) \times E^1$ and $M_2 \cong E^* \times U(3)$ are the two maximal Levi subgroups.

We consider representations of the Levi subgroups, extend them trivially to the unipotent subgroups to obtain representations of the parabolic groups. We now perform normalised parabolic induction to obtain representations of $U(5)$.

In this article we give a classification of the irreducible unitary representations of $U(5)$ in terms of Langlands quotients. At first we consider the irreducible subquotients obtained by induction from representations of M_0 and from non-cuspidal, not fully-induced representations of M_1 and M_2 (the subquotients determined in [Sch14]). We then consider the irreducible subquotients of representations induced from cuspidal representations of $M_1 \cong \mathrm{GL}(2,E) \times E^1$.

2. Definitions

Let G be a connected reductive algebraic group, defined over a non-archimedean local field of characteristic 0. Let V be a vector space, defined over the complex numbers. Let π be a representation of G on V. We denote it by (π, V) and sometimes by π or V.

Let $\mathcal{K}$ denote the set of open compact subgroups of G .

Definition 2.1 A representation (π, V) of G is said to be **smooth** if every $v \in V$ is fixed by a neighbourhood of the unity in G . This is equivalent to saying that $\exists K \in \mathcal{K}$ such that $\pi(k)v = v \; \forall k \in K$.

Definition 2.2 A representation (π, V) of G is said to be **admissible** if it is smooth and for every $K \in \mathcal{K}$ the space V^K of fixed vectors under K is finite dimensional.

From now on let (π, V) be an admissible representation of G .

Definition 2.3 Let (π, V) be a smooth representation of G . Let $V^* = \mathrm{Hom}(V, \mathbb{C})$ be the space of linear forms on V. One defines a representation (π^*, V^*) of G : if $v^* \in V^*$ and $g \in G$ then $\pi^*(g)(v^*)$ is defined by $v \mapsto v^*(\pi(g^{-1})(v))$. Let $\widetilde{V}$ be the subspace of smooth vectors in V^*, i.e. the subset $\widetilde{V} \subset V^*$ of elements $\widetilde{v}$ such that the stabiliser of $\widetilde{v}$ is an open subgroup of G . One shows that $\widetilde{V}$ is invariant under G for the action of π^*. So π^* induces a representation $\widetilde{\pi}$ on $\widetilde{V}$; $(\widetilde{\pi}, \widetilde{V})$ is called the **dual representation** of (π, V).

Definition 2.4 A representation π is called **hermitian** if $\pi \cong \overline{\widetilde{\pi}}$.

Definition 2.5 A representation (π, V) of a group G is called a **unitary representation** if and only if on the vector space V there exists a positive definite hermitean form $\langle \ , \ \rangle : V \times V \to \mathbb{C}$ that is invariant under the action of G :

$$\langle \pi(g)v, \pi(g)w \rangle = \langle v, w \rangle \; \forall g \in \mathrm{G}, \forall v, w \in V.$$

Definition 2.6 A **matrix coefficient** $f_{v,\widetilde{v}}$ of a representation (π, V) is a (locally constant) function: $f_{v,\widetilde{v}} : \mathrm{G} \to \mathbb{C} : g \mapsto \widetilde{v}(\pi(g)(v))$, where $v \in V$ and $\widetilde{v} \in \widetilde{V}$.

Definition 2.7 An irreducible representation π of G is called **cuspidal** if π has a non-zero matrix coefficient $f : \mathrm{G} \to \mathbb{C}$ that is compactly supported modulo the center of G.

Definition 2.8 An irreducible representation π of G is called **square-integrable** if π is unitary and if π has a non-zero matrix coefficient $f : \mathrm{G} \to \mathbb{C}$ that is square-integrable modulo the center Z of G : $f \in L^2(\mathrm{G}/Z)$. It follows that all matrix coefficients of π are square-integrable.

Definition 2.9 An irreducible representation π of a group G is **tempered** if it is unitary and if π has a non-zero matrix coefficient $f : \mathrm{G} \to \mathbb{C}$ that verifies $f \in L^{2+\epsilon}(\mathrm{G}/Z)\ \forall \epsilon > 0$.

Remark 2.10 Square-integrable representations are tempered.

Let F be a non-archimedean local field of characteristic 0. i.e. $\mathbb{Q}_p$ or a finite extension of $\mathbb{Q}_p$, where p is a prime number.

Let $E : F$ be a field extension of degree two, hence a Galois extension. Let $\mathrm{Gal}(E : F) = \{\mathrm{id}, \sigma\}$. The action of the non-trivial element σ of the Galois group is called the conjugation of elements in E corresponding to the extension $E : F$. We write $\sigma(x) = \overline{x}\ \forall\ x \in E$ and extend $^-$ to matrices with entries in E.

Let $\Phi \in \mathrm{GL}(n, E)$ be a hermitian matrix, i.e. $\overline{\Phi}^t = \Phi$, let U_Φ be the unitary group definded by Φ :

$$U_\Phi = \{g \in \mathrm{GL}(n, E) : g\Phi\overline{g}^t = \Phi\}.$$

Let $\Phi_n = (\Phi_{ij})$, where $\Phi_{ij} = (-1)^{i-1}\delta_{i,n+1-j}$ and δ_{ab} is the Cronecker delta.

Let $\zeta \in E^*$ be an element of trace 0, i.e. $tr(\zeta) = \zeta + \overline{\zeta} = 0$.

If n is odd, then $\Phi_n = \begin{pmatrix} & & & & 1 \\ & & & \iddots & \\ & & 1 & & \\ & -1 & & & \\ 1 & & & & \end{pmatrix}$ is hermitian. If n is even, $\zeta\Phi_n = \begin{pmatrix} & & & & \zeta \\ & & & \iddots & \\ & & -\zeta & & \\ & \zeta & & & \\ -\zeta & & & & \end{pmatrix}$ is hermitian.

Denote by $U(n)$ the **unitary group** corresponding to Φ_n if n is odd or to $\zeta\Phi_n$ if n is even, respectively. It is quasi split.

Let n be an even positive integer. We will call Levi subgroup of $U(n)$ a subgroup of block diagonal matrices

$$L := \{\begin{pmatrix} A_1 & & & & & & & - & 0 \\ & A_2 & & & & & & & | \\ & & \ddots & & & & & & \\ & & & A_k & & & & & \\ & & & & B & & & & \\ & & & & & {}^t\overline{A}_k^{-1} & & & \\ & & & & & & \ddots & & \\ | & & & & & & & & \\ 0 & - & & & & & & {}^t\overline{A}_1^{-1} & \end{pmatrix},\ A_i \in \mathrm{GL}_{n_i}(E)\ \text{ for }\ 1 \le i \le k \text{ and } B \in U(m)\},$$

where $m, n_1, \ldots, n_k$ are strictly positive integers such that $m + 2\sum_{i=1}^{k} n_i = n$. (If $k = 0$, then there are no n_i and $L = U(n)$.)

It is canonically isomorphic to the product $\mathrm{GL}(n_1,E)\times\ldots\times\mathrm{GL}(n_k,E)\times U(m)$. We chose the corresponding parabolic subgroup P such that it contains L and the subgroup of upper triangular matrices in $U(n)$. We call a parabolic subgroup P that contains the subgroup of upper triangular matrices standard. Let N be the unipotent subgroup with identity matrices for the block diagonal matrices of L, arbitray entries in E above and $0's$ below. Then one has the Levi decomposition $P=LN$.

Let $\pi_i,\ i=1,\ldots k$, be smooth admissible representations of $\mathrm{GL}(n_i,E)$ and σ a smooth admissible representation of $U(m)$. Let $\pi_1\otimes\ldots\otimes\pi_k\otimes\sigma$ denote the representation of $L=\mathrm{GL}(n_1,E)\times\ldots\times\mathrm{GL}(n_k,E)\times U(m)$ and denote by $\pi:=\mathrm{Ind}_P^{U(n)}(\pi_1\otimes\ldots\pi_k\otimes\sigma)=\pi_1\times\ldots\pi_k\rtimes\sigma$ the normalized parabolically induced representation, where P is the corresponding standard parabolic subgroup containing L.

Definition 2.11 Let π be an irreducible representation of $\mathrm{GL}(n,E)$. Then there exist irreducible cuspidal representations $\rho_1,\rho_2,\ldots,\rho_k$ of general linear groups that are, up to isomorphism, uniquely defined by π, such that π is isomorphic to a subquotient of $\rho_1\times\ldots\times\rho_k$. The multiset of equivalence classess $(\rho_1,\ldots,\rho_k)$ is called the **cuspidal support** of π. It is denoted by $\mathrm{supp}(\pi)$.

Definition 2.12 Let $n\in\mathbb{N}$ and let τ be an irreducible representation of $U(n)$. Then there exist irreducible cuspidal representations $\rho_1,\ldots,\rho_k$ of general linear groups and an irreducible cuspidal representation σ of some $U(m)$ that are, up to isomorphism and replacement of ρ_i by $\rho_i^{-1}(-)$ for some $i\in\{1,\ldots,k\}$, uniquely defined by τ, s. t. τ is isomorphic to a subquotient of $\rho_1\times\ldots\times\rho_k\rtimes\sigma$. The representation σ is called the **partial cuspidal support** of τ and is denoted by τ_{cusp}.

Definition 2.13 Let π be a smooth representation of finite length of G. Then $\hat{\pi}$ denotes the **Aubert dual** of π, as defined in [Aub95].

3. Irreducible unitary representations of $U(5)$, in terms of Langlands-quotients

3.1. **Representations with cuspidal support in M_0, fully-induced.** Let $\chi_{\omega_{E/F}}\in X_{\omega_{E/F}}$. Let $\pi_{1,\chi_{\omega_{E/F}}}$ be the unique irreducible square-integrable subquotient of $|\ |^{1/2}\ \chi_{\omega_{E/F}}\rtimes\lambda'$. Let $\chi_{1_{F^*}}\in X_{1_{F^*}}$. Recall that $\chi_{1_{F^*}}=\sigma_{1,\chi_{1_{F^*}}}\oplus\sigma_{2,\chi_{1_{F^*}}}$, where $\sigma_{1,\chi_{1_{F^*}}}$ and $\sigma_{2,\chi_{1_{F^*}}}$ are tempered [Key84].

Proposition 3.1. *Let $0<\alpha_2\leq\alpha_1,\alpha>0$. Let χ_1,χ_2 and χ be unitary characters of E^*. The following list exhausts all irreducible hermitian representations of $U(5)$ with cuspidal support in M_0 :*

0. tempered representations of $U(5)$,
1. $\mathrm{Lg}(|\ |^{\alpha_1}\chi_1;|\ |^{\alpha_2}\chi_2;\lambda')$ where $\chi_1,\chi_2\in X_{N_{E/F}(E^*)}$ or $\alpha_1=\alpha_2$ and $\chi_1(x)=\chi_2^{-1}(\overline{x})\ \forall x\in E^*$,
2. $\mathrm{Lg}(|\ |^{\alpha}\chi_1;\chi_2\rtimes\lambda')$ where $\chi_1\in X_{N_{E/F}(E^*)}$ and $\chi_2\notin X_{1_{F^*}}$,
3. $\mathrm{Lg}(|\ |^{\alpha}\chi\,\mathrm{St}_{\mathrm{GL}_2};\lambda')$ where $\chi\in X_{N_{E/F}(E^*)}$,
4. $\mathrm{Lg}(|\ |^{\alpha}\chi;\lambda'(\det)\,\mathrm{St}_{U(3)}),\mathrm{Lg}(|\ |^{\alpha}\chi,\pi_{1,\chi_{\omega_{E/F}}}),\mathrm{Lg}(|\ |^{\alpha}\chi;\sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^{\alpha}\chi;\sigma_{2,\chi_{1_{F^*}}})$, where $\chi\in X_{N_{E/F}(E^*)}$.

Outline of the proof: 0. Tempered representations are unitary, hence hermitian.

1.-4. Let $\lambda_i,\ i=0,1,2$ be representations of the Levi subgroups M_0, M_1 and M_2. By [Cas95], $\widetilde{\mathrm{Ind}_{M_i}^{U(5)}}(\lambda_i) \cong \mathrm{Ind}_{M_i}^{U(5)}(\lambda_i),\ i=0,1,2$, is equivalent to $\exists w \in W$ such that $\overline{\widetilde{\lambda_i}} = w\lambda_i,\ i=0,1,2$. This holds also for the Langlands quotients, and the proof is immediate.

Irreducible subquotients of $\chi_1 \times \chi_2 \rtimes \lambda'$

Let χ_1, χ_2 be unitary characters of E^*.

All irreducible subquotients of $\chi_1 \times \chi_2 \rtimes \lambda'$ are tempered, hence unitary.

$$\mathrm{Lg}(|\ |^{\alpha_1}\chi_1; |\ |^{\alpha_2}\chi_2; \lambda'),\ 0<\alpha_2\le\alpha_1,\ \chi_1 \notin X_{N_{E/F}(E^*)} \text{ or } \chi_2 \notin X_{N_{E/F}(E^*)}$$

Theorem 3.2. *Let χ_1, χ_2 be unitary characters of E^* such that $\chi_1 \notin X_{N_{E/F}(E^*)}$ or $\chi_2 \notin X_{N_{E/F}(E^*)}$.*

1. Let $0<\alpha_2\le\alpha_1$. Let $\alpha_1 \neq \alpha_2$ or $\exists x \in E^*$ s. t. $\chi_1(x) \neq \chi_2^{-1}(\overline{x})$. $\mathrm{Lg}(|\ |^{\alpha_1}\chi_1; |\ |^{\alpha_2}\chi_2; \lambda')$ is non-unitary.

2. Let $0<\alpha_1=\alpha_2$ and $\chi_1(x) = \chi_2^{-1}(\overline{x})\ \forall x \in E^*$. $\mathrm{Lg}(|\ |^{\alpha_1}\chi_1; |\ |^{\alpha_2}\chi_2; \lambda')$ is unitary for $0<\alpha_1=\alpha_2\le 1/2$.

$\mathrm{Lg}(|\ |^{\alpha_1}\chi_1; |\ |^{\alpha_2}\chi_2; \lambda')$ is non-unitary for $\alpha > 1/2$.

Proof. 1. Let $\alpha_1 \neq \alpha_2$ or $\exists x \in E^*$ such that $\chi_1(x) \neq \chi_2^{-1}(\overline{x})$. The representations $|\ |^{\alpha_1}\chi_1 \times |\ |^{\alpha_2}\chi_2 \rtimes \lambda'$ are not hermitian, by [Cas95] 3.1.2, $\mathrm{Lg}(|\ |^{\alpha_1}\chi_1; |\ |^{\alpha_2}\chi_2; \lambda')$ is not hermitian, hence not unitary.

2. Let $\alpha_1=\alpha_2$ and $\chi_1(x) = \chi_2^{-1}(\overline{x})\ \forall x \in E^*$. Representations $|\ |^{\alpha_1}\chi_1 \times |\ |^{\alpha_2}\chi_2 \rtimes \lambda'$ are hermitian. Let $\alpha_1=\alpha_2<1/2$ and $\chi_1(x) = \chi_2^{-1}(\overline{x})\ \forall x \in E^*$. Representations $|\ |^{\alpha_1}\chi_1 \times |\ |^{\alpha_2}\chi_2 \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.3] and equal to their Langlands quotients $\mathrm{Lg}(|\ |^{\alpha_1}\chi_1; |\ |^{\alpha_2}\chi_2; \lambda')$. $\chi_1 \times \chi_2 \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.1] and unitary. For $\alpha_1=\alpha_2<1/2$, representations $\mathrm{Lg}(|\ |^{\alpha_1}\chi_1; |\ |^{\alpha_2}\chi_2; \lambda')$ form a continuous 1-parameter family of irreducible hermitian representations that we realize on the same vector space V (for a detailed version of this argument in a similar case see the proof of Theorem 3.4). By Lemma [Sch14, Lemma 3.6] $\mathrm{Lg}(|\ |^{\alpha_1}\chi_1; |\ |^{\alpha_2}\chi_2; \lambda')$ is unitary for $\alpha_1=\alpha_2<1/2$. For $\alpha_1=\alpha_2=1/2$ and $\chi_1(x) = \chi_2^{-1}(\overline{x})\ \forall x \in E^*$, $|\ |^{1/2}\chi_1 \times |\ |^{1/2}\chi_2 \rtimes \lambda'$ is reducible by [Sch14, Th. 4.3]. By [Mil73] $\mathrm{Lg}(|\ |^{1/2}\chi_1; |\ |^{1/2}\chi_2; \lambda')$ is unitary.

Let $\alpha_1=\alpha_2>1/2$ and $\chi_1(x) = \chi_2^{-1}(\overline{x})\ \forall x \in E^*$. Representations $|\ |^{\alpha_1}\chi_1 \times |\ |^{\alpha_2}\chi_2 \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.3] and equal to their Langlands-quotients $\mathrm{Lg}(|\ |^{\alpha_1}\chi_1; |\ |^{\alpha_2}\chi_2; \lambda')$. By [Sch14, Lemmas 3.6 and 3.8] $\mathrm{Lg}(|\ |^{\alpha_1}\chi_1; |\ |^{\alpha_2}\chi_2; \lambda')$ is non-unitary for $\alpha_1=\alpha_2>1/2$. □

$$\mathrm{Lg}(|\ |^{\alpha}\chi_1; \chi_2 \rtimes \lambda'),\ \alpha>0,\ \chi_1 \notin X_{N_{E/F}(E^*)} \textbf{ or } \chi_2 \notin X_{N_{E/F}(E^*)}$$

Let $\chi_{1_{F^*}} \in X_{1_{F^*}}$. Recall that $\chi_{1_{F^*}} \rtimes \lambda' = \sigma_{1,\chi_{1_{F^*}}} \oplus \sigma_{2,\chi_{1_{F^*}}}$, where $\sigma_{1,\chi_{1_{F^*}}}$ and $\sigma_{2,\chi_{1_{F^*}}}$ are tempered [Key84].

Recall that $X_{N_{E/F}(E^*)} = 1 \cup X_{\omega_{E/F}} \cup X_{1_{F^*}}$.

Theorem 3.3. *1. Let $\alpha > 0$. Let $\chi_1 \notin X_{N_{E/F}(E^*)}$ and $\chi_2 \notin X_{1_{F^*}}$. $\mathrm{Lg}(|\ |^\alpha \chi_1; \chi_2 \rtimes \lambda')$ is non-unitary.*

2. Let $\alpha > 0$. Let $\chi_1 \notin X_{N_{E/F}(E^*)}$. $\mathrm{Lg}(|\ |^\alpha \chi_1; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^\alpha \chi_1; \sigma_{2,\chi_{1_{F^*}}})$ are non-unitary.

3. Let $\alpha > 0$. Let $\chi_1 \in X_{N_{E/F}(E^*)}$ and $\chi_2 \notin X_{N_{E/F}(E^*)}$.

3.1 Let $\chi_1 = 1$. Let $0 < \alpha \leq 1$. $\mathrm{Lg}(|\ |^\alpha 1; \chi_2 \rtimes \lambda')$ is unitary.

Let $\alpha > 1$. $\mathrm{Lg}(|\ |^\alpha 1; \chi_2 \rtimes \lambda')$ is non-unitary.

3.2 Let $\chi_1 \in X_{\omega_{E/F}}$. Let $0 < \alpha \leq 1/2$. $\mathrm{Lg}(|\ |^\alpha \chi_{\omega_{E/F}}; \chi_2 \rtimes \lambda')$ is unitary.

Let $\alpha > 1/2$. $\mathrm{Lg}(|\ |^\alpha \chi_{\omega_{E/F}}; \chi_2 \rtimes \lambda')$ is non-unitary.

3.3 Let $\chi_1 \in X_{1_{F^*}}$. Let $\alpha > 0$. $\mathrm{Lg}(|\ |^\alpha \chi_{1_{F^*}}; \chi_2 \rtimes \lambda')$ is non-unitary.

Proof. 1. For $\alpha > 0$, representations $|\ |^\alpha \chi_1 \times \chi_2 \rtimes \lambda'$ are not hermitian. By [Cas95] 3.1.2, $\mathrm{Lg}(|\ |^\alpha \chi_1; \chi_2 \rtimes \lambda')$ is not hermitian, hence not unitary.

2. For $\alpha > 0$, $|\ |^\alpha \chi_1 \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $|\ |^\alpha \chi_1 \rtimes \sigma_{2,\chi_{1_{F^*}}}$ are not hermitian. By [Cas95] $\mathrm{Lg}(|\ |^\alpha \chi_1; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^\alpha \chi_1; \sigma_{2,\chi_{1_{F^*}}})$ are not hermitian and hence non-unitary.

3.1 $1 \times \chi_2 \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.1] and unitary. For $0 < \alpha < 1$, representations $|\ |^\alpha 1 \times \chi_2 \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.5] and equal to their Langlands-quotient $\mathrm{Lg}(|\ |^\alpha 1; \chi_2 \rtimes \lambda')$. By [Sch14, Lemma 3.6] these Langlands-quotients are unitary. For $\alpha = 1$, $|\ | 1 \times \chi_2 \rtimes \lambda'$ reduces for the first time, by [Sch14, Th. 4.5]. By [Mil73] $\mathrm{Lg}(|\ |\ 1; \chi_2 \rtimes \lambda')$ is unitary. For $\alpha > 1$, representations $|\ |^\alpha 1 \times \chi_2 \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.5] and equal to their Langlands-quotient $\mathrm{Lg}(|\ |^\alpha 1; \chi_2 \rtimes \lambda')$. By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary.

3.2 $\chi_{\omega_{E/F}} \times \chi_2 \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.1] and unitary. For $0 < \alpha < 1/2$, representations $|\ |^\alpha \chi_{\omega_{E/F}} \times \chi_2 \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.5] and equal to their Langlands-quotient $\mathrm{Lg}(|\ |^\alpha \chi_{\omega_{E/F}}; \chi_2 \rtimes \lambda')$. By [Sch14, Lemma 3.6] these Langlands-quotients are unitary. For $\alpha = 1/2$, $|\ |^{1/2} \chi_{\omega_{E/F}} \times \chi_2 \rtimes \lambda'$ reduces for the first time, by [Sch14, Th. 4.5]. By [Mil73] $\mathrm{Lg}(|\ |^{1/2} \chi_{\omega_{E/F}}; \chi_2 \rtimes \lambda')$ is unitary. For $\alpha > 1/2$, representations $|\ |^\alpha \chi_{\omega_{E/F}} \times \chi_2 \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.5] and equal to their Langlands-quotient $\mathrm{Lg}(|\ |^\alpha \chi_{\omega_{E/F}}; \chi_2 \rtimes \lambda')$. By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary.

3.3 For $\alpha > 0$, representations $|\ |^\alpha \chi_{1_{F^*}} \times \chi_2 \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.5] and equal to their Langlands-quotient $\mathrm{Lg}(|\ |^\alpha \chi_{1_{F^*}}; \chi_2 \rtimes \lambda')$. By [Sch14, Lemma 3.6] and [Sch14, Lemma 3.8] these Langlands-quotients are non-unitary. □

We now take $\chi_1, \chi_2 \in X_{N_{E/F}(E^*)}$.

$$\mathrm{Lg}(|\ |^{\alpha_1} 1; |\ |^{\alpha_2} 1; \lambda'),\ 0 < \alpha_2 \leq \alpha_1,\ \mathrm{Lg}(|\ |^\alpha 1; 1 \rtimes \lambda'),\ \alpha > 0$$

Theorem 3.4. *1. Let $0 < \alpha_2 \leq \alpha_1 \leq 1$ and $\alpha_1 + \alpha_2 \leq 1$. $\mathrm{Lg}(|\ |^{\alpha_1} 1; |\ |^{\alpha_2} 1; \lambda')$ is unitary.*

2. Let $0 < \alpha_2 \leq \alpha_1$, $\alpha_1 + \alpha_2 > 1$, $(\alpha_1, \alpha_2) \neq (2, 1)$. If $\alpha_1 = 1$, then let $\alpha_2 \notin (0, 1]$. Then $\mathrm{Lg}(|\ |^{\alpha_1} 1; |\ |^{\alpha_2} 1; \lambda')$ is non-unitary.

2.1 $\mathrm{Lg}(|\ |^2 1; |\ | 1; \lambda') = 1_{U(5)}$ is unitary.

3. For $0 < \alpha \leq 1$, $\mathrm{Lg}(|\ |^\alpha 1; 1 \rtimes \lambda')$ is unitary.

4. For $\alpha > 1$, $\mathrm{Lg}(|\ |^\alpha 1; 1 \rtimes \lambda')$ is non-unitary.

Proof. 1. $1 \times 1 \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.1] and unitary. Let $0 < \alpha_2 \leq \alpha_1$. For $\alpha_1 + \alpha_2 < 1$, representations $|\ |^{\alpha_1} 1\times |\ |^{\alpha_2} 1 \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.3], hence equal to $\mathrm{Lg}(|\ |^{\alpha_1} 1; |\ |^{\alpha_2} 1; \lambda')$.

We will now construct a continuous one-parameter family of hermitian representations. Let $0 < \alpha_2 \leq \alpha_1$ such that $\alpha_1 + \alpha_2 \leq 1$. Let π_{α_1,α_2} denote the two-parameter family of hermitian representations $|\ |^{\alpha_1} 1\times |\ |^{\alpha_2} 1 \rtimes \lambda'$. Let V_{α_1,α_2} be the vector space of π_{α_1,α_2}.

Recall the Levi decomposition $P_0 = M_0 N_0$, where P_0 is the minimal parabolic subgroup of $U(5)$, $M_0 = \{\begin{pmatrix} x & & & & 0 \\ & y & & & \\ & & k & & \\ & & & \overline{y}^{-1} & \\ 0 & & & & \overline{x}^{-1} \end{pmatrix},\ x, y \in E^*,\ k \in E^1\}$ is the minimal Levi subgroup and N_0 the unipotent radical of P_0.

Let (π, V) be the extension to P_0 of the representation $1 \otimes 1 \otimes \lambda'$ of M_0. Let δ_{P_0} denote the modulus character of P_0.

$V_{0,0} = \{f : \mathrm{G} \to V : f \quad \text{is smooth and } f(pg) = \delta_{P_0}(p)\pi(p)f(g) \forall g \in \mathrm{G}\}$.

$V_{\alpha_1,\alpha_2} = \{h : \mathrm{G} \to V : h \quad \text{is smooth and } h(pg) = \delta_{P_0}(p) \mid x \mid^{\alpha_1} \mid y \mid^{\alpha_2} \pi(p)h(g) \forall g \in \mathrm{G}\}$,

where $p \in P_0$, $p = \begin{pmatrix} x & & - & & * \\ & y & & & | \\ & & k & & \\ & | & & \overline{y}^{-1} & \\ 0 & - & & & \overline{x}^{-1} \end{pmatrix}$, $x, y \in E^*, k \in E^1, * \in E$. Let $\mathcal{O}$ denote the ring of integers of E. $|\ |: E^* \to F^*$ is non-ramified, hence $(x, y) \mapsto |\ x\ |^{\alpha_1} |\ y\ |^{\alpha_2}$, $x, y \in E^*$, is trivial on $E^1 \times E^1 \cong \mathcal{O}^* \times \mathcal{O}^*$. Let $K := U(\mathcal{O})$, it is a maximal compact subgroup of G . We have $\mathrm{G} = KP_0$. Let $f \in V_{0,0}$. There exists a unique extension of $f_{|K} : \mathrm{G} \to V$ to a function $h \in V_{\alpha_1,\alpha_2}$, so $f_{|K} = h_{|K}$. This induces an isomorphism $T_{\alpha_1,\alpha_2} : V_{0,0} \xrightarrow{\sim} V_{\alpha_1,\alpha_2}$. Via the composition with T_{α_1,α_2} we consider all representations π_{α_1,α_2} in $V_{0,0}$.

Let $w \in W$ be the longest element of the Weyl group. Let

$A(w, \lambda) :|\ |^{\alpha_1} 1\times |\ |^{\alpha_2} 1 \rtimes \lambda' \to |\ |^{-\alpha_1} 1\times |\ |^{-\alpha_2} 1 \rtimes \lambda'$ be the standard long intertwining operator.

On $V_{0,0}$ we define a set of non-degenerate hermitian forms $\langle\ ,\ \rangle_{\alpha_1,\alpha_2}$ by

$$\langle f, h\rangle_{\alpha_1,\alpha_2} = \int_{U(\mathcal{O})} A(w, \lambda) f(k)\overline{h(k)}dk,\ f, h \in V_{0,0},$$

such that $\langle\ ,\ \rangle_{\alpha_1,\alpha_2}$ is invariant by $T^{-1}_{\alpha_1,\alpha_2}\pi_{\alpha_1,\alpha_2}T_{\alpha_1,\alpha_2}$.

Fix α_1 and α_2 such that $\alpha_1 + \alpha_2 = 1$. Let $\pi_t = \pi_{t\alpha_1,t\alpha_2}$, $t \in [0, 1]$ denote a continuous one-parameter family of hermitian representations. Let V_t be the vector space of π_t. Via the isomorphism $T_t : V_0 \xrightarrow{\sim} V_t$, we consider all representations π_t in V_0, as before.

Choose a polynomial $p(t)$ with real coefficents, such that $A(t) = p(t)A(w, \lambda)$ is holomorphic and non-zero for $t \in [0, 1]$. So for the one-parameter family of representations π_t one obtains, on the same space V_0, a set of non-degenerate hermitian forms $\langle\ ,\ \rangle_t$ given by

$$\langle f, h\rangle_t = \int_{U(\mathcal{O})} A(t) f(k)\overline{h(k)}dk,\ f, h \in V_0,$$

such that $\langle\ ,\ \rangle_t$ is invariant under $T_t^{-1}\pi_t T_t$.

$\langle\ ,\ \rangle_0$ is positive definite, hence by [Sch14, Lemma 3.6] $\langle\ ,\ \rangle_t$ is positive definite until $|\ |^{t\alpha_1} 1\times |\ |^{t\alpha_2} \rtimes \lambda'$ reduces for the first time, for $t = 1$. By [Mil73], for $t = 1$, the irreducible subquotients of

$|\,|^{\alpha_1}\,1\times|\,|^{\alpha_2}\,1\rtimes\lambda'$ are unitary. Hence for $0<\alpha_2\leq\alpha_1\leq 1,\ \alpha_1+\alpha_2\leq 1$, the Langlands quotients $\mathrm{Lg}(|\,|^{\alpha_1}\,1;|\,|^{\alpha_2}\,1;\lambda')$ are unitary.

2. $+$ 2.1 $|\,|^2\,1\times|\,|\,1\rtimes\lambda'$ is reducible. $\lambda'(\det)\,\mathrm{St}_{U(3)}$ is the unique irreducible square-integrable subquotient of $|\,|\,1\rtimes\lambda'$ [Key84]. By [Cas81, p.915] the subquotient $|\,|^{3/2}\,1_{\mathrm{GL}_2}\rtimes\lambda'$ is reducible and has the subquotients $1_{U(5)}=\mathrm{Lg}(|\,|^2\,1;|\,|\,1;\lambda')$ and $\mathrm{Lg}(|\,|^2\,1;\lambda'(\det)\,\mathrm{St}_{U(3)})$. $1_{U(5)}=\mathrm{Lg}(|\,|^2\,1;|\,|\,1;\lambda')$ is unitary, $\mathrm{Lg}(|\,|^2\,1;\lambda'(\det)\,\mathrm{St}_{U(3)})$ is non-unitary ([Cas81]). For $1/2<\alpha<3/2$, representations $|\,|^{\alpha}\,1_{\mathrm{GL}_2}\rtimes\lambda'$ are irreducible by [Sch14, Th. 4.7], they form a continuous one-parameter family of irreducible hermitian representations on the space V_α. Like before we identify the vectorspaces V_α for $1/2<\alpha<3/2$. For $\alpha=3/2$, the irreducible subquotient $\mathrm{Lg}(|\,|^2\,1;\lambda'(\det)\,\mathrm{St}_{U(3)})$ of $|\,|^{3/2}\,1_{\mathrm{GL}_2}\rtimes\lambda'$ is not unitary [Cas81]. Hence, by [Mil73] and by [Sch14, Lemma 3.6], $|\,|^{\alpha}\,1_{\mathrm{GL}_2}\rtimes\lambda'=\mathrm{Lg}(|\,|^{\alpha_1}\,1;|\,|^{\alpha_2}\,1;\lambda')$ is non-unitary for $1/2<\alpha<3/2$, i.e. for $1<\alpha_1<2,\ \alpha_1-\alpha_2=1$.

By [Cas81, p.915] the subquotient $|\,|^2\,1\rtimes\lambda'(\det)1_{U(3)}$ of the representation $|\,|^2\,1\times|\,|\,1\rtimes\lambda'$ is reducible. It has the subquotients $1_{U(5)}=\mathrm{Lg}(|\,|^2\,1;|\,|\,1;\lambda')$ and $\mathrm{Lg}(|\,|^{3/2}\,\mathrm{St}_{\mathrm{GL}_2};\lambda')$. $1_{U(5)}$ is unitary, $\mathrm{Lg}(|\,|^{3/2}\,\mathrm{St}_{\mathrm{GL}_2};\lambda')$ is non-unitary [Cas81].

Let $1<\alpha<2$. Representations $|\,|^{\alpha}\,1\rtimes\lambda'(\det)1_{U(3)}$ are irreducible by [Sch14, Th. 4.7], they form a continuous one-parameter family of irreducible hermitian representations on the space V_α. Similar as before we identify V_α for $1<\alpha<2$. The irreducible subquotient $\mathrm{Lg}(|\,|^{3/2}\,\mathrm{St}_{\mathrm{GL}_2};\lambda')$ of $|\,|^2\,1\rtimes\lambda'(\det)1_{U(3)}$ is non-unitary [Cas81]. Hence by [Mil73] and by [Sch14, Lemma 3.6] representations $|\,|^{\alpha}\,1\rtimes\lambda'(\det)1_{U(3)}=\mathrm{Lg}(|\,|^{\alpha_1}\,1;|\,|^1\,1;\lambda')$ are non-unitary for $1=\alpha_2<\alpha_1=\alpha<2$.

Let $1<\alpha_1<2,\ 0<\alpha_2<1,\ \alpha_1-\alpha_2<1$. $|\,|^{\alpha_1}\,1\times|\,|^{\alpha_2}\,1\rtimes\lambda'$ is irreducible by [Sch14, Th. 4.3] and equal to its own Langlands quotient $\mathrm{Lg}(|\,|^{\alpha_1}\,1;|\,|^{\alpha_2}\,1;\lambda')$. Fix $0<\alpha_2<1$ and let $1<\alpha_1\leq\alpha_2+1$. Let π_{α_1} denote the continuous one-parameter family of hermitian representations $|\,|^{\alpha_1}\,1\times|\,|^{\alpha_2}\,1\rtimes\lambda'$ on the same vector space V. For $\alpha_1=\alpha_2+1$ irreducible subquotients of the representations $|\,|^{\alpha_1}\,1\times|\,|^{\alpha_2}\,1\rtimes\lambda'$ are non-unitary, as seen in the previous paragraph. By [Mil73] and by [Sch14, Lemma 3.6] the Langlands quotients $\mathrm{Lg}(|\,|^{\alpha_1}\,1;|\,|^{\alpha_2}\,1;\lambda')$ are non-unitary. (II in Figure 1 on page 9)

Let $\alpha_1>2,\ \alpha_1-\alpha_2=1$. $\mathrm{Lg}(|\,|^{\alpha_1}\,1;|\,|^{\alpha_2}\,1;\lambda')$ is non-unitary by Lemmas **??** and **??**. For $\alpha_1>2,\ \alpha_2=1$, $\mathrm{Lg}(|\,|^{\alpha_1}\,1;|\,|^{\alpha_2}\,1;\lambda')$ is non-unitary by the same argument.

The same holds for $\mathrm{Lg}(|\,|^{\alpha_1}\,1;|\,|^{\alpha_2}\,1;\lambda')$, where $\alpha_1>1,0<\alpha_2<1,\ \alpha_1-\alpha_2>1$, for $1<\alpha_2\leq\alpha_1,\ \alpha_1-\alpha_2<1$ and for $\alpha_1>2,\ \alpha_1-\alpha_2>1$ (III, IV, V in Figure **??** on page **??**).

Let $\alpha_1=1,\ \alpha_2\in(0,1]$. We have no proof that $\mathrm{Lg}(|\,|\,1;|\,|^{\alpha_2}\,1;\lambda')$ is non-unitary.

3. $1\times1\times\lambda'$ is irreducible by Theorem [Sch14, Th. 4.1] and unitary. For $0<\alpha<1$, $|\,|^{\alpha}\,1\times1\rtimes\lambda'$ is irreducible by [Sch14, Th. 4.5] and equal to its own Langlands quotient $\mathrm{Lg}(|\,|^{\alpha}\,1;1\rtimes\lambda')$. By Lemma [Sch14, Lemma 3.6] these representations are unitary. For $\alpha=1$, $|\,|\,1\times1\rtimes\lambda'$ reduces for the first time, hence $\mathrm{Lg}(|\,|\,1;1\rtimes\lambda')$ is unitary [Mil73].

4. For $\alpha>1$, $|\,|^{\alpha}\,1\times1\rtimes\lambda'=\mathrm{Lg}(|\,|^{\alpha}\,1;1\rtimes\lambda')$ is irreducible by [Sch14, Th. 4.5] and by [Sch14, Lemmas 3.6 and 3.8] non-unitary. □

$$\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F}};|\,|^{\alpha_2}\,\chi_{\omega_{E/F}};\lambda'),\ 0<\alpha_2\leq\alpha_1,\ \mathrm{Lg}(|\,|^{\alpha}\,\chi_{\omega_{E/F}};\chi_{\omega_{E/F}}\rtimes\lambda'),\ \alpha>0,\ \ \chi_{\omega_{E/F}}\in X_{\omega_{E/F}}$$

Let $\chi_{\omega_{E/F}}\in X_{\omega_{E/F}}$.




Figure 1

FIGURE 1. Let $\alpha_1, \alpha_2 \geq 0$. Figure 1 shows lines and points of reducibility of the representation $|\ |^{\alpha_1} 1\times |\ |^{\alpha_2} 1 \rtimes \lambda'$ and the unitary dual. Let $0 < \alpha_2 \leq \alpha_1$. $\mathrm{Lg}(|\ |^{\alpha_1} 1; |\ |^{\alpha_2} 1; \lambda')$ is unitary for $0 < \alpha_2 \leq \alpha_1 < 1, \alpha_1 + \alpha_2 \leq 1$ and for $\alpha_1 = 2,\ \alpha_2 = 1$. Except for $\alpha_1 = 1,\ 0 < \alpha_2 \leq 1$, it is non-unitary. $\mathrm{Lg}(|\ |^{\alpha} 1; 1 \rtimes \lambda')$ is unitary for $0 < \alpha \leq 1$ and non-unitary for $\alpha > 1$.

Theorem 3.5. *1. Let* $0 < \alpha_2 \leq \alpha_1 \leq 1/2$. $\mathrm{Lg}(|\ |^{\alpha_1} \chi_{\omega_{E/F}}; |\ |^{\alpha_2} \chi_{\omega_{E/F}}; \lambda')$ *is unitary.*

2. Let $\alpha_1 > 1/2,\ \alpha_2 \leq \alpha_1,\ (\alpha_1, \alpha_2) \neq (3/2, 1/2)$. *If* $0 < \alpha_2 < 1/2$, *then let* $\alpha_1 \notin (1/2, 1-\alpha_2]$. $\mathrm{Lg}(|\ |^{\alpha_1} \chi_{\omega_{E/F}}; |\ |^{\alpha_2} \chi_{\omega_{E/F}}; \lambda')$ *is non-unitary.*

3. Let $0 < \alpha \leq 1/2$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}}; \chi_{\omega_{E/F}} \rtimes \lambda')$ *is unitary.*

4. Let $\alpha > 1$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}}; \chi_{\omega_{E/F}} \rtimes \lambda')$ *is non-unitary.*

Proof. 1. $\chi_{\omega_{E/F}} \times \chi_{\omega_{E/F}} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.1] and unitary. For $0 < \alpha_2 \leq \alpha_1 < 1/2$ $|\ |^{\alpha_1} \chi_{\omega_{E/F}} \times |\ |^{\alpha_2} \chi_{\omega_{E/F}} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.3] and equal to its Langlands quotient $\mathrm{Lg}(|\ |^{\alpha_1} \chi_{\omega_{E/F}}; |\ |^{\alpha_2} \chi_{\omega_{E/F}}; \lambda')$. Let $\alpha_1 = 1/2$ and fix $0 < \alpha_2 \leq 1/2$. For $t \in [0,1]$, let $\pi_{(t1/2, t\alpha_2)} =: \pi_t$ denote the continuous one-parameter family of hermitian representations $|\ |^{t1/2} \chi_{\omega_{E/F}} \times |\ |^{t\alpha_2} \chi_{\omega_{E/F}} \rtimes \lambda'$. For $t \in [0,1)$ these representations are equal to their own Langlands quotient $\mathrm{Lg}(|\ |^{t1/2} \chi_{\omega_{E/F}}; |\ |^{t\alpha_2} \chi_{\omega_{E/F}}; \lambda')$ and by [Sch14, Lemma 3.6] unitary. For $t = 1$ the representations π_t reduce for the first time. By [Mil73] $\mathrm{Lg}(|\ |^{1/2} \chi_{\omega_{E/F}}; |\ |^{\alpha_2} \chi_{\omega_{E/F}}; \lambda')$ is unitary.

2. Let $1 < \alpha_1 < 3/2$ and let $\alpha_1 - \alpha_2 = 1$. Then $|\ |^{\alpha_1} \chi_{\omega_{E/F}} \times |\ |^{\alpha_2} \chi_{\omega_{E/F}} \rtimes \lambda'$ is reducible by [Sch14, Th. 4.3]. The subquotients $|\ |^{\frac{\alpha_1+\alpha_2}{2}} \chi_{\omega_{E/F}} 1_{\mathrm{GL}_2} \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.7]. They form a continuous 1-parameter family of irreducible hermitian representations, that similar as before, we realize on the same vector space V. Let $\alpha_1 = 3/2,\ \alpha_2 = 1/2$. $|\ |^{\frac{3/2+1/2}{2}} \chi_{\omega_{E/F}} 1_{\mathrm{GL}_2} \rtimes \lambda' = |\ |\ \chi_{\omega_{E/F}} 1_{\mathrm{GL}_2} \rtimes \lambda'$ reduces by [Sch14, Th. 5.4]. Let $\pi_{1,\chi_{\omega_{E/F}}}$ be the unique square-integrable subquotient of $|\ |^{1/2} \chi_{\omega_{E/F}} \rtimes \lambda'$ [Key84]. By Theorem 1.1 and Remark 4.7 in [HT10] the irreducible subquotient $\mathrm{Lg}(|\ |^{3/2} \chi_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}})$ of $|\ |\ \chi_{\omega_{E/F}} 1_{\mathrm{GL}_2} \rtimes \lambda'$ is non-unitary. By [Mil73] and [Sch14, Lemma 3.6] the representations $|\ |^{\frac{\alpha_1+\alpha_2}{2}} \chi_{\omega_{E/F}} 1_{\mathrm{GL}_2} \rtimes \lambda',\ 1 < \alpha_1 < 3/2,\ \alpha_1 - \alpha_2 = 1$, that are equal to $\mathrm{Lg}(|\ |^{\alpha_1} \chi_{\omega_{E/F}}; |\ |^{\alpha_2} \chi_{\omega_{E/F}}; \lambda'),\ 1 < \alpha_1 < 3/2,\ \alpha_1 - \alpha_2 = 1$, are non-unitary.

Let $1/2 < \alpha_1 < 3/2,\ \alpha_2 = 1/2$. The representations $|\ |^{\alpha_1} \chi_{\omega_{E/F}} \times |\ |^{1/2} \chi_{\omega_{E/F}} \rtimes \lambda'$ are reducible by [Sch14, Th. 4.3]. Let $\pi_{2,\chi_{\omega_{E/F}}}$ be the unique irreducible non-tempered subquotient of $|\ |^{1/2}$

$\chi_{\omega_{E/F}} \rtimes \lambda'$ [Key84]. For $1/2 < \alpha_1 < 3/2$, the representations $|\ |^{\alpha_1}\ \chi_{\omega_{E/F}} \rtimes \pi_{2,\chi_{\omega_{E/F}}}$ are irreducible by [Sch14, Th. 4.11] and equal to the Langlands quotient $\mathrm{Lg}(|\ |^{\alpha_1}\ \chi_{\omega_{E/F}}; |\ |^{1/2}\ \chi_{\omega_{E/F}}; \lambda')$. They form a 1 - parameter family of irreducible hermitian representations, that we realise on the same vector space V. For $\alpha_1 = 3/2$, $|\ |^{3/2}\ \chi_{\omega_{E/F}} \rtimes \pi_{2,\chi_{\omega_{E/F}}}$ reduces by [Sch14, Th. 5.4], and by Theorem 1.1 and Remark 4.7 in [HT10] its irreducible subquotient $\mathrm{Lg}(|\ |\ \chi_{\omega_{E/F}}\ \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is non-unitary. By [Mil73] and [Sch14, Lemma 3.6] $|\ |^{\alpha_1}\ \chi_{\omega_{E/F}} \rtimes \pi_2 = \mathrm{Lg}(|\ |^{\alpha_1}\ \chi_{\omega_{E/F}}; |\ |^{1/2}\ \chi_{\omega_{E/F}}; \lambda')$ is non-unitary for $1/2 < \alpha_1 < 3/2$.

Representations $|\ |^{\alpha_1}\ \chi_{\omega_{E/F}} \times |\ |^{\alpha_2}\ \chi_{\omega_{E/F}} \rtimes \lambda'$ in II,III,IV,V of Figure 2 on page 11 are irreducible by [Sch14, Th. 4.3] and equal to their own Langlands quotient $\mathrm{Lg}(|\ |^{\alpha_1}\ \chi_{\omega_{E/F}}; |\ |^{\alpha_2}\ \chi_{\omega_{E/F}}; \lambda')$. The Langlands quotients $\mathrm{Lg}(|\ |^{\alpha_1}\ \chi_{\omega_{E/F}}; |\ |^{\alpha_2}\ \chi_{\omega_{E/F}}; \lambda')$ in II are non-unitary by [Mil73] and [Sch14, Lemma 3.6]. $\mathrm{Lg}(|\ |^{\alpha_1}\ \chi_{\omega_{E/F}}; |\ |^{\alpha_2}\ \chi_{\omega_{E/F}}; \lambda')$ in III, IV and V are non-unitary by [Sch14, Lemmas 3.6 and 3.8].

3. $\chi_{\omega_{E/F}} \times \chi_{\omega_{E/F}} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.1] and unitary. For $0 < \alpha < 1/2$ $|\ |^{\alpha}\ \chi_{\omega_{E/F}} \times \chi_{\omega_{E/F}} \rtimes \lambda'$ is irreducible by Theorem [Sch14, Th. 4.5] and equal to its Langlands quotient $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \chi_{\omega_{E/F}} \rtimes \lambda')$. By [Sch14, Lemma 3.6] these Langlands quotients are unitary.

For $\alpha = 1/2$, $|\ |^{1/2}\ \chi_{\omega_{E/F}} \times \chi_{\omega_{E/F}} \rtimes \lambda'$ reduces for the first time ([Sch14, Th. 4.5]). By [Mil73] $\mathrm{Lg}(|\ |^{1/2}\ \chi_{\omega_{E/F}}; \chi_{\omega_{E/F}} \rtimes \lambda')$ is unitary.

4. For $\alpha > 1$, $|\ |^{\alpha}\ \chi_{\omega_{E/F}} \times \chi_{\omega_{E/F}} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.5] and equal to its Langlands quotient $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \chi_{\omega_{E/F}}; \lambda')$. By [Sch14, Lemmas 3.6 and 3.8] these Langlands quotients are non-unitary. □

Remark 3.6 We do not have a proof that the representation $\mathrm{Lg}(|\ |^{3/2}\ \chi_{\omega_{E/F}} \times |\ |^{1/2}\ \chi_{\omega_{E/F}} \rtimes \lambda')$ is unitary. It is the Aubert dual of a square-integrable representation and should be unitary, see [Sch14, Th. 5.4]. See [Han09], where the proof is given for orthogonal and symplectic groups.

Remark 3.7 In the Grothendieck group of the category of admissible representations of finite length one has $|\ |\ \chi_{\omega_{E/F}} \times \chi_{\omega_{E/F}} \rtimes \lambda' = |\ |^{1/2}\ \chi_{\omega_{E/F}}\ \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda' + |\ |^{1/2}\ \chi_{\omega_{E/F}} 1_{\mathrm{GL}_2} \rtimes \lambda'$. If we assume that $|\ |^{1/2}\ \chi_{\omega_{E/F}}\ \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ and $|\ |^{1/2}\ \chi_{\omega_{E/F}} 1_{\mathrm{GL}_2} \rtimes \lambda'$ are irreducible (see [Sch14, Remark 5.5]), we are able to prove that $\mathrm{Lg}(|\ |^{\alpha_1}\ \chi_{\omega_{E/F}}; |\ |^{\alpha_2}\ \chi_{\omega_{E/F}}; \lambda')$ is non-unitary for $1/2 < \alpha_1 < 1,\ \alpha_2 \leq 1 - \alpha_1$ and that $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \chi_{\omega_{E/F}}; \lambda')$ is non-unitary for $1/2 < \alpha \leq 1$:

Let $1/2 < \alpha_1 < 1,\ \alpha_2 = 1 - \alpha_1$. By [Sch14, Th. 4.3] representations $|\ |^{\alpha_1}\ \chi_{\omega_{E/F}} \times |\ |^{\alpha_2}\ \chi_{\omega_{E/F}} \rtimes \lambda'$ are reducible. By [Sch14, Th. 4.7] the subquotient $|\ |^{\frac{\alpha_1 - \alpha_2}{2}}\ \chi_{\omega_{E/F}} 1_{\mathrm{GL}_2} \rtimes \lambda'$ is irreducible. It is equal to $\mathrm{Lg}(|\ |^{\alpha_1}\ \chi_{\omega_{E/F}}; |\ |^{\alpha_2}\ \chi_{\omega_{E/F}}; \lambda')$. By assumption $|\ |^{1/2}\ \chi_{\omega_{E/F}} 1_{\mathrm{GL}_2} \rtimes \lambda'$ is irreducible, it is equal to $\mathrm{Lg}(|\ |\ \chi_{\omega_{E/F}}; \chi_{\omega_{E/F}} \rtimes \lambda')$. Hence we can extend the argument 2 in the proof of 3.5: $\mathrm{Lg}(|\ |^{\alpha_1}\ \chi_{\omega_{E/F}}; |\ |^{\alpha_2}\ \chi_{\omega_{E/F}}; \lambda')$ is non-unitary for $1/2 < \alpha_1 < 1$ and $\alpha_2 = 1 - \alpha_1$ and for $1 < \alpha_1 < 3/2,\ \alpha_2 = \alpha_1 - 1$, and $\mathrm{Lg}(|\ |\ \chi_{\omega_{E/F}}; \chi_{\omega_{E/F}} \rtimes \lambda')$ is non-unitary.

Let $1/2 < \alpha_1 < 1,\ \alpha_2 < 1 - \alpha_1$. By [Sch14, Th. 4.3] the representations $|\ |^{\alpha_1}\ \chi_{\omega_{E/F}} \times |\ |^{\alpha_2}\ \chi_{\omega_{E/F}} \rtimes \lambda'$ are irreducible, they are equal to their Langlands-quotient $\mathrm{Lg}(|\ |^{\alpha_1}\ \chi_{\omega_{E/F}}; |\ |^{\alpha_2}\ \chi_{\omega_{E/F}}; \lambda')$ (I in Figure 2, page 11). By [Mil73] and by [Sch14, Lemma 3.6] these Langlands-quotients are non-unitary.

Let $1/2 < \alpha < 1$. By [Sch14, Th. 4.5] representations $|\ |^{\alpha}\ \chi_{\omega_{E/F}} \times \chi_{\omega_{E/F}} \rtimes \lambda'$ are irreducible, they are equal to their Langlands-quotient $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \chi_{\omega_{E/F}} \rtimes \lambda')$. $|\ |\ \chi_{\omega_{E/F}} \times \chi_{\omega_{E/F}} \rtimes \lambda'$ is

reducible by [Sch14, Th. 4.5], $\mathrm{Lg}(|\ |\chi_{\omega_{E/F}} \times \chi_{\omega_{E/F}}; \lambda')$ is non-unitary by the foregoing argument. By [Mil73] and by [Sch14, Lemma 3.6] $\mathrm{Lg}(|\ |^{\alpha}\chi_{\omega_{E/F}}; \chi_{\omega_{E/F}} \rtimes \lambda')$ is non-unitary for $1/2 < \alpha < 1$.




Figure 2

FIGURE 2. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$. Figure 2 shows lines and points of reducibility of the representation $|\ |^{\alpha_1}\chi_{\omega_{E/F}} \times |\ |^{\alpha_2}\chi_{\omega_{E/F}} \rtimes \lambda'$ and the unitary dual. Let $0 < \alpha_2 \leq \alpha_1$. $\mathrm{Lg}(|\ |^{\alpha_1}\chi_{\omega_{E/F}}; |\ |^{\alpha_2}\chi_{\omega_{E/F}}; \lambda')$ is unitary for $0 < \alpha_2 \leq \alpha_1 \leq 1/2$. Except for $1/2 < \alpha_1 < 1$, $0 < \alpha_2 \leq 1 - \alpha_1$ and for $\alpha_1 = 3/2$, $\alpha_2 = 1/2$, it is non-unitary. $\mathrm{Lg}(|\ |^{\alpha}\chi_{\omega_{E/F}}; \chi_{\omega_{E/F}} \rtimes \lambda')$ is unitary for $0 < \alpha \leq 1/2$. For $\alpha > 1$ it is non-unitary.

$$\mathrm{Lg}(|\ |^{\alpha_1}\chi_{1_{F^*}}; |\ |^{\alpha_2}\chi_{1_{F^*}}; \lambda'), 0 < \alpha_2 \leq \alpha_1$$

Theorem 3.8. *Let* $\chi_{1_{F^*}} \in X_{1_{F^*}}$.

1. *Let* $0 < \alpha_2 \leq \alpha_1$, $\alpha_1 + \alpha_2 > 1$. *Then* $\mathrm{Lg}(|\ |^{\alpha_1}\chi_{1_{F^*}}; |\ |^{\alpha_2}\chi_{1_{F^*}}; \lambda')$ *is non-unitary.*
2. *Let* $0 < \alpha_2 \leq \alpha_1$, $\alpha_1 + \alpha_2 = 1$. *Then* $\mathrm{Lg}(|\ |^{\alpha_1}\chi_{1_{F^*}}; |\ |^{\alpha_2}\chi_{1_{F^*}}; \lambda')$ *is unitary.*

Proof. 1. Let $\alpha_1 - \alpha_2 = 1$. Then $|\ |^{\alpha_1}\chi_{1_{F^*}} \times |\ |^{\alpha_2}\chi_{1_{F^*}} \rtimes \lambda'$ is reducible by [Sch14, Th. 4.3]. Let $\alpha := \frac{\alpha_1+\alpha_2}{2}$, then $\alpha > 1/2$. The subquotient $|\ |^{\alpha}\chi_{1_{F^*}}1_{\mathrm{GL}_2} \rtimes \lambda'$ of $|\ |^{\alpha_1}\chi_{1_{F^*}} \times |\ |^{\alpha_2}\chi_{1_{F^*}} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.7] and equal to the Langlands quotient $\mathrm{Lg}(|\ |^{\alpha_1}\chi_{1_{F^*}}; |\ |^{\alpha_2}\chi_{1_{F^*}}; \lambda')$. Representations $|\ |^{\alpha}\chi_{1_{F^*}}1_{\mathrm{GL}_2} \rtimes \lambda'$ form a continuous 1 - parameter family of irreducible hermitian representations that we realise on the same vector space V. If there existed $\alpha > 1/2$ such that $|\ |^{\alpha}\chi_{1_{F^*}}1_{\mathrm{GL}_2} \rtimes \lambda'$ was unitary, then by [Sch14, Lemma 3.6] $|\ |^{\alpha}\chi_{1_{F^*}}1_{\mathrm{GL}_2} \rtimes \lambda'$ would be unitary $\forall \alpha > 1/2$, in contradiction to [Sch14, Lemma 3.8].

The representations $|\ |^{\alpha_1}\chi_{1_{F^*}} \times |\ |^{\alpha_2}\chi_{1_{F^*}} \rtimes \lambda'$ in II and III of Figure 3, page 12, are irreducible by [Sch14, Th. 4.3] and equal to their own Langlands quotient $\mathrm{Lg}(|\ |^{\alpha_1}\chi_{1_{F^*}}; |\ |^{\alpha_2}\chi_{1_{F^*}}; \lambda')$. Similar as in Theorem 3.4 (2) we show that $\mathrm{Lg}(|\ |^{\alpha_1}\chi_{1_{F^*}}; |\ |^{\alpha_2}\chi_{1_{F^*}}; \lambda')$ is non-unitary if $\alpha_1 + \alpha_2 > 1$ and $\alpha_1 - \alpha_2 < 1$, and if $\alpha_1 + \alpha_2 > 1$ and $\alpha_1 - \alpha_2 > 1$ ([Sch14, Lemmas 3.6 and 3.8]).

2. $|\ |^{\alpha_1}\chi_{1_{F^*}} \times |\ |^{\alpha_2}\chi_{1_{F^*}} \rtimes \lambda'$ is reducible by [Sch14, Th. 4.3]. Let $\alpha := \frac{\alpha_1-\alpha_2}{2}$. The subquotient $|\ |^{\alpha}\chi_{1_{F^*}}1_{\mathrm{GL}_2} \rtimes \lambda'$ of $|\ |^{\alpha_1}\chi_{1_{F^*}} \times |\ |^{\alpha_2}\chi_{1_{F^*}} \rtimes \lambda'$ is irreducible by [Sch14, Th. 6.7] and equal to




Figure 3

FIGURE 3. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{1_{F^*}} \in X_{1_{F^*}}$. Figure 3 shows lines and points of reducibility of the representation $|\ |^{\alpha_1} \chi_{1_{F^*}} \times |\ |^{\alpha_2} \chi_{1_{F^*}} \rtimes \lambda'$ and the unitary dual. Let $0 < \alpha_2 \leq \alpha_1$. $\mathrm{Lg}(|\ |^{\alpha_1} \chi_{1_{F^*}}; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda')$ is unitary for $\alpha_1 + \alpha_2 = 1$. It is non-unitary for $\alpha_1 + \alpha_2 > 1$.

the Langlands quotient $\mathrm{Lg}(|\ |^{\alpha_1} \chi_{1_{F^*}}; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda')$. These representations form a continuous 1 - parameter family of irreducible hermitian representations that we realise on the same vector space V. $\chi_{1_{F^*}} 1_{\mathrm{GL}_2} \rtimes \lambda'$ is irreducible by [Sch14, Prop. 4.8]. It is unitary. For $\alpha = 1/2$, $|\ |^{1/2} \chi_{1_{F^*}} 1_{\mathrm{GL}_2} \rtimes \lambda'$ reduces for the first time ([Sch14, Th. 4.5]). By [Sch14, 3.6] the representations $|\ |^{\alpha} \chi_{1_{F^*}} 1_{\mathrm{GL}_2} \rtimes \lambda' = \mathrm{Lg}(|\ |^{\alpha_1} \chi_{1_{F^*}}; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda')$ are unitary for $0 \leq \alpha < 1/2$, i.e. for $0 < \alpha_2 \leq \alpha_1,\ \alpha_1 + \alpha_2 = 1$. □

Remark 3.9 Let $0 < \alpha_2 \leq \alpha_1,\ \alpha_1 + \alpha_2 < 1$. We do not have a proof that $\mathrm{Lg}(|\ |^{\alpha_1} \chi_{1_{F^*}}; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda')$ is non-unitary.

In the following Theorems 3.10, 3.11 and 3.12, when speaking of the Langlands-quotient, we will exceptionally allow that $\alpha_1 < \alpha_2$ for ease of notation.

$$\mathrm{Lg}(|\ |^{\alpha_1} 1; |\ |^{\alpha_2} \chi_{\omega_{E/F}}; \lambda'),\ \alpha_1, \alpha_2 > 0,\ \mathrm{Lg}(|\ |^{\alpha} 1; \chi_{\omega_{E/F}} \rtimes \lambda'),\ \mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}}; 1 \rtimes \lambda'),\ \alpha > 0,\ \chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$$

Let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$. Let $\pi_{1,\chi_{\omega_{E/F}}}$ denote the unique square-integrable subquotient and let $\pi_{2,\chi_{\omega_{E/F}}}$ denote the unique non-tempered subquotient of $|\ |^{1/2} \chi_{\omega_{E/F}} \rtimes \lambda'$.

Theorem 3.10. *Let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$.*

1. *Let $0 < \alpha_1 \leq 1,\ 0 < \alpha_2 \leq 1/2$. $\mathrm{Lg}(|\ |^{\alpha_1} 1; |\ |^{\alpha_2} \chi_{\omega_{E/F}}; \lambda')$ is unitary.*
2. *Let $\alpha_1 > 1,\ \alpha_2 > 0$, or let $0 < \alpha_1 \leq 1,\ \alpha_2 > 1/2$. $\mathrm{Lg}(|\ |^{\alpha_1} 1; |\ |^{\alpha_2} \chi_{\omega_{E/F}}; \lambda')$ is non-unitary.*
3. *Let $0 < \alpha \leq 1$. $\mathrm{Lg}(|\ |^{\alpha} 1; \chi_{\omega_{E/F}} \rtimes \lambda')$ is unitary.*
4. *Let $\alpha > 1$. $\mathrm{Lg}(|\ |^{\alpha} 1; \chi_{\omega_{E/F}} \rtimes \lambda')$ is non-unitary.*
5. *Let $\alpha \leq 1/2$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}}; 1 \rtimes \lambda')$ is unitary.*
6. *Let $\alpha > 1/2$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}}; 1 \rtimes \lambda')$ is non-unitary.*

Proof. 1. $1 \times \chi_{\omega_{E/F}} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.1] and unitary. $|\ |^{\alpha_1} 1 \times |\ |^{\alpha_2} \chi_{\omega_{E/F}} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.3] and equal to its own Langlands-quotient $\mathrm{Lg}(|\ |^{\alpha_1} 1 \times |\ |^{\alpha_2} \chi_{\omega_{E/F}} \rtimes \lambda')$.

Like in 3.4 we construct continuous one-parameter families of irreducible hermitian representations and obtain that $\mathrm{Lg}(|\ |^{\alpha_1} 1\times |\ |^{\alpha_2}\ \chi_{\omega_{E/F}} \rtimes \lambda')$ is unitary for $0<\alpha_1<1,\ 0<\alpha_2<1/2$ ([Sch14, Lemma 3.6]).

For $\alpha_1=1,\ 0<\alpha_2\leq 1/2,\ |\ |1\times |\ |^{\alpha_2}\ \chi_{\omega_{E/F}} \rtimes \lambda'$ reduces for the first time ([Sch14, Th. 4.3]). For $0<\alpha_1<1,\ \alpha_2=1/2,\ |\ |^{\alpha_1} 1\times |\ |^{1/2}\ \chi_{\omega_{E/F}} \rtimes \lambda'$ reduces for the first time ([Sch14, Th. 4.3]). By [Mil73] $\mathrm{Lg}(|\ |^{\alpha_1} 1\times |\ |^{\alpha_2}\ \chi_{\omega_{E/F}} \rtimes \lambda')$ is unitary for $\alpha_1=1,\ 0<\alpha_2\leq 1/2$, and for $0<\alpha_1<1,\ \alpha_2=1/2$.

2. Let $\alpha_2=1/2$. Let $\alpha>1$. The representation $|\ |^{\alpha} 1\times |\ |^{1/2}\ \chi_{\omega_{E/F}} \rtimes \lambda'$ is reducible by [Sch14, Th. 4.3]. Its subquotient $|\ |^{\alpha} 1 \rtimes \pi_{2,\chi_{\omega_{E/F}}}$ is irreducible by [Sch14, Th. 4.11] and equal to $\mathrm{Lg}(|\ |^{\alpha} 1;|\ |^{1/2}\ \chi_{\omega_{E/F}};\lambda')$. The argument is similar to 3.4; if there existed $\alpha>1$, such that $\mathrm{Lg}(|\ |^{\alpha} 1;|\ |^{1/2}\ \chi_{\omega_{E/F}};\lambda')$ was unitary, then by [Sch14, Lemma 3.6] $\mathrm{Lg}(|\ |^{\alpha} 1;|\ |^{1/2}\ \chi_{\omega_{E/F}};\lambda')$ would be unitary $\forall\alpha>1$, in contradiction to [Sch14, Lemma 6.8].

Let $\alpha_1=1$. Let $\alpha>1/2$. The representation $|\ |1\times |\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \lambda'$ is reducible by [Sch14, Th. 4.3]. The subquotient $|\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \lambda'(\det)1_{U(3)}$ is irreducible by [Sch14, Th. 4.9] and equal to $\mathrm{Lg}(|\ |1;|\ |^{\alpha}\ \chi_{\omega_{E/F}};\lambda')$. By [Sch14, Lemmas 3.6 and 3.8] $\mathrm{Lg}(|\ |1;|\ |^{\alpha}\ \chi_{\omega_{E/F}};\lambda')$ is non-unitary for $\alpha>1/2$.

Let $\alpha_1>1,\ 1/2\neq\alpha_2>0$, or let $0<\alpha_1<1,\ \alpha_2>1/2$. Representations $|\ |^{\alpha_1} 1\times |\ |^{\alpha_2}\ \chi_{\omega_{E/F}} \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.3] and equal to their own Langlands-quotient $\mathrm{Lg}(|\ |^{\alpha_1} 1;|\ |^{\alpha_2}\ \chi_{\omega_{E/F}};\lambda')$. Like in 3.4 we obtain that these Langlands-quotients are non-unitary (II, III and IV in Figure 4, page 14).

3 +4. $1\times\chi_{\omega_{E/F}} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.1] and unitary. $|\ |^{\alpha} 1\times\chi_{\omega_{E/F}} \rtimes \lambda'$ is irreducible for $0<\alpha<1$ by [Sch14, Th. 4.5] and equal to its own Langlands -quotient $\mathrm{Lg}(|\ |^{\alpha} 1;\chi_{\omega_{E/F}} \rtimes \lambda')$. By [Sch14, Lemma 3.6] these subquotients are unitary. The representation $|\ |1\times\chi_{\omega_{E/F}} \rtimes \lambda'$ is reducible by [Sch14, Th. 4.5]. The subquotient $\chi_{\omega_{E/F}} \rtimes \lambda'(\det)1_{U(3)}$ is irreducible by [Sch14, Prop. 4.10] and equal to $\mathrm{Lg}(|\ |1;\chi_{\omega_{E/F}} \rtimes \lambda')$. It is unitary. By [Sch14, Th. 4.5] representations $|\ |^{\alpha} 1\times\chi_{\omega_{E/F}} \rtimes \lambda'$ are irreducible for $\alpha>1$ and equal to their own Langlands quotient $\mathrm{Lg}(|\ |^{\alpha} 1;\chi_{\omega_{E/F}} \rtimes \lambda')$. By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary.

5+6. $1\times\chi_{\omega_{E/F}} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.1] and unitary. By [Sch14, Th. 4.5] $1\times |\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \lambda'$ is irreducible for $0<\alpha<1/2$ and equal to its own Langlands -quotient $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}};1 \rtimes \lambda')$. By [Sch14, Lemma 3.6] these subquotients are unitary. By [Sch14, Th. 4.5] the representation $1\times |\ |^{1/2}\ \chi_{\omega_{E/F}} \rtimes \lambda'$ is reducible. Its subquotient $1 \rtimes \pi_{2,\chi_{\omega_{E/F}}}$ is irreducible by [Sch14, Prop. 4.12] and equal to the Langlands-quotient $\mathrm{Lg}(|\ |^{1/2}\ \chi_{\omega_{E/F}},1 \rtimes \lambda')$. It is unitary. Representations $1\times |\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \lambda'$ are irreducible for $\alpha>1/2$ by [Sch14, Th. 4.5] and equal to their own Langlands quotient $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}};1 \rtimes \lambda')$. By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary. □

$$\mathrm{Lg}(|\ |^{\alpha_1} 1;|\ |^{\alpha_2}\ \chi_{1_{F^*}};\lambda'),\ \alpha_1,\alpha_2>0,\ \mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}};1 \rtimes \lambda'),\ \alpha>0,\ \chi_{1_{F^*}}\in X_{1_{F^*}}$$

Let $\chi_{1_{F^*}}\in X_{1_{F^*}}$.

Theorem 3.11. *1. Let $\alpha_1,\alpha_2>0$. $\mathrm{Lg}(|\ |^{\alpha_1} 1;|\ |^{\alpha_2}\ \chi_{1_{F^*}};\lambda')$ is non-unitary.*

2. Let $\alpha>0$. $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}};1 \rtimes \lambda')$ is non-unitary.




Figure 4

FIGURE 4. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$. Figure 4 shows lines and points of reducibility of the representation $|\ |^{\alpha_1} 1\times |\ |^{\alpha_2} \chi_{\omega_{E/F}} \rtimes \lambda'$ and the unitary dual. $\mathrm{Lg}(|\ |^{\alpha_1} 1; |\ |^{\alpha_2} \chi_{\omega_{E/F}}; \lambda')$ is unitary for $0 < \alpha_1 \leq 1,\ 0 < \alpha_2 \leq 1/2$. Otherwise it is non-unitary. $\mathrm{Lg}(|\ |^{\alpha} 1;\ \chi_{\omega_{E/F}} \rtimes \lambda')$ is unitary for $0 < \alpha \leq 1$, $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}}; 1 \rtimes \lambda')$ is unitary for $0 < \alpha \leq 1/2$. Otherwise these Langlands-quotients are non-unitary.

Proof. 1. Let $\alpha_1 = 1, \alpha_2 > 0$. The representation $|\ |\ 1\times |\ |^{\alpha_2} \chi_{1_{F^*}} \rtimes \lambda'$ is reducible by [Sch14, Th. 4.3]. Its subquotient $|\ |^{\alpha_2} \chi_{1_{F^*}} \rtimes \lambda'(\det)1_{U(3)}$ is irreducible by [Sch14, Th. 4.9] and equal to $\mathrm{Lg}(|\ |\ 1; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda')$. By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary.

Let $\alpha_1, \alpha_2 > 0$, $\alpha_1 \neq 1$. Representations $|\ |^{\alpha_1} 1\times |\ |^{\alpha_2} \chi_{1_{F^*}} \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.3] and equal to their own Langlands-quotient $\mathrm{Lg}(|\ |^{\alpha_1} 1; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda')$. Similar as in 3.4 (2), by [Sch14, Lemmas 3.6 and 3.8], we find that these Langlands-quotients are non-unitary.

2. Let $\alpha > 0$. Representations $|\ |^{\alpha} \chi_{1_{F^*}} \times 1 \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.5] and equal to their own Langlands-quotient $\mathrm{Lg}(|\ |^{\alpha} \chi_{1_{F^*}}; 1 \rtimes \lambda')$. By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary. (Figure 5, page 15)

□

$$\mathrm{Lg}(|\ |^{\alpha_1} \chi_{\omega_{E/F}}; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda'),\ \alpha_1,\ \alpha_2 > 0,\ \mathrm{Lg}(|\ |^{\alpha} \chi_{1_{F^*}}; \chi_{\omega_{E/F}} \rtimes \lambda'),\ \alpha > 0,\ \chi_{\omega_{E/F}} \in X_{\omega_{E/F}},\ \chi_{1_{F^*}} \in X_{1_{F^*}}$$

Let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$, let $\chi_{1_{F^*}} \in X_{1_{F^*}}$.

Theorem 3.12. *1. Let $\alpha_1, \alpha_2 > 0$. $\mathrm{Lg}(|\ |^{\alpha_1} \chi_{\omega_{E/F}}; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda')$ is non-unitary.*
2. Let $\alpha > 0$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{1_{F^*}}; \chi_{\omega_{E/F}} \rtimes \lambda')$ is non-unitary.

Proof. 1. Let $\alpha_1 = 1/2, \alpha_2 > 0$. The representation $|\ |^{1/2} \chi_{\omega_{E/F}}\times |\ |^{\alpha_2} \chi_{1_{F^*}} \rtimes \lambda'$ is reducible by [Sch14, Th. 4.3]. Let $\pi_{2,\chi_{\omega_{E/F}}}$ be the unique irreducible non-tempered subquotient of $|\ |^{1/2} \chi_{\omega_{E/F}} \rtimes \lambda'$. The subquotient $|\ |^{\alpha_2} \chi_{1_{F^*}} \rtimes \pi_{2,\chi_{\omega_{E/F}}}$ of $|\ |^{1/2} \chi_{\omega_{E/F}}\times |\ |^{\alpha_2} \chi_{1_{F^*}} \rtimes \lambda'$ is irreducible




Figure 5

FIGURE 5. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{1_{F^*}} \in X_{1_{F^*}}$. Figure 5 shows lines and points of reducibility of the representation $|\ |^{\alpha_1} 1 \times |\ |^{\alpha_2} \chi_{1_{F^*}} \rtimes \lambda'$ and the unitary dual. $\mathrm{Lg}(|\ |^{\alpha_1} 1; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda')$ is non-unitary $\forall \alpha_1, \alpha_2 > 0$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{1_{F^*}}; 1 \rtimes \lambda')$ is non-unitary $\forall \alpha > 0$.

by [Sch14, Th. 4.11] and equal to $\mathrm{Lg}(|\ |^{1/2} \chi_{\omega_{E/F}}; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda')$. By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary.

Let $\alpha_1, \alpha_2 > 0$, $\alpha_1 \neq 1/2$. Representations $|\ |^{\alpha_1} \chi_{\omega_{E/F}} \times |\ |^{\alpha_2} \chi_{1_{F^*}} \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.3] and equal to their own Langlands-quotient $\mathrm{Lg}(|\ |^{\alpha_1} \chi_{\omega_{E/F}}; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda')$. Similar as in 3.4 (2), by [Sch14, Lemmas 3.6 and 3.8], we find that these Langlands-quotients are non-unitary.

2. Let $\alpha > 0$. Representations $|\ |^{\alpha} \chi_{1_{F^*}} \times \chi_{\omega_{E/F}} \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.5] and equal to their own Langlands-quotient $\mathrm{Lg}(|\ |^{\alpha} \chi_{1_{F^*}}; \chi_{\omega_{E/F}} \rtimes \lambda')$. By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary. (Figure 6, page 15) □




Figure 6

FIGURE 6. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}, \chi_{1_{F^*}} \in X_{1_{F^*}}$. Figure 6 shows lines and points of reducibility of the representation $|\ |^{\alpha_1} \chi_{\omega_{E/F}} \times |\ |^{\alpha_2} \chi_{1_{F^*}} \rtimes \lambda'$ and the unitary dual. $\mathrm{Lg}(|\ |^{\alpha_1} \chi_{\omega_{E/F}}; |\ |^{\alpha_2} \chi_{1_{F^*}}; \lambda')$ is non-unitary $\forall \alpha_1, \alpha_2 > 0$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{1_{F^*}}; \chi_{\omega_{E/F}} \rtimes \lambda')$ is non-unitary $\forall \alpha > 0$.

$$\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2};\lambda'),\ 0<\alpha_2\leq\alpha_1,\ \mathrm{Lg}(|\,|^{\alpha}\,\chi_{\omega_{E/F},1};\chi_{\omega_{E/F},2}\rtimes\lambda'),\alpha>0,\ \chi_{\omega_{E/F},1},\chi_{\omega_{E/F},2}\in X_{\omega_{E/F}},\ \chi_{\omega_{E/F},1}\not\cong\chi_{\omega_{E/F},2}$$

Theorem 3.13. *Let $\chi_{\omega_{E/F},1},\chi_{\omega_{E/F},2}\in X_{\omega_{E/F}}$, such that $\chi_{\omega_{E/F},1}\not\cong\chi_{\omega_{E/F},2}$.*

1. *Let $0<\alpha_2\leq\alpha_1\leq 1/2$. $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2};\lambda')$ is unitary.*
2. *Let $\alpha_1>1/2$, $0<\alpha_2\leq\alpha_1$. $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2}\rtimes\lambda')$ is non-unitary.*
3. *Let $0<\alpha\leq 1/2$. $\mathrm{Lg}(|\,|^{\alpha}\,\chi_{\omega_{E/F},1};\chi_{\omega_{E/F},2}\rtimes\lambda')$ is unitary.*
4. *Let $\alpha>1/2$. $\mathrm{Lg}(|\,|^{\alpha}\,\chi_{\omega_{E/F},1};\chi_{\omega_{E/F},2}\rtimes\lambda')$ is non-unitary.*

Proof. 1. $\chi_{\omega_{E/F},1}\times\chi_{\omega_{E/F},2}\rtimes\lambda'$ is irreducible by [Sch14, Th. 4.1] and unitary. Let $\alpha_1=1/2$, fix $0<\alpha_2\leq 1/2$ and let $t\in[0,1]$. Representations $|\,|^{t1/2}\,\chi_{\omega_{E/F},1}\times|\,|^{t\alpha_2}\,\chi_{\omega_{E/F},2}\rtimes\lambda'$ give a continuous 1- parameter family of hermitian representations that we realise on the same vector space V, similar as in 3.4. For $t\in[0,1)$, $|\,|^{t1/2}\,\chi_{\omega_{E/F},1}\times|\,|^{t\alpha_2}\,\chi_{\omega_{E/F},2}\rtimes\lambda'$ is irreducible by [Sch14, Th. 4.3] and unitary by [Sch14, Lemma 3.6]. For $t=1$, $|\,|^{1/2}\,\chi_{\omega_{E/F},1}\times|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2}\rtimes\lambda'$ reduces for the first time. By [Mil73] $\mathrm{Lg}(|\,|^{1/2}\,\chi_{\omega_{E/F},1};|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2};\lambda')$ is unitary. Hence $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2};\lambda')$ is unitary for $0<\alpha_2\leq\alpha_1\leq 1/2$.

2. Let $\alpha_1>1/2$, $\alpha_2=1/2$. $|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1}\times|\,|^{1/2}\,\chi_{\omega_{E/F},2}\rtimes\lambda'$ is reducible by [Sch14, Th. 4.3]. Let $\pi_{2,\chi'_{\omega_{E/F}}}$ be the unique irreducible non-tempered subquotient of $|\,|^{1/2}\,\chi_{\omega_{E/F},2}\rtimes\lambda'$. By [Sch14, Th. 4.11] $|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1}\rtimes\pi_{2,\chi'_{\omega_{E/F}}}=\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{1/2}\,\chi_{\omega_{E/F},2};\lambda')$ is irreducible. For $\alpha_1>1/2$, $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{1/2}\,\chi_{\omega_{E/F},2};\lambda')$ gives a continuous 1-parameter-family of irreducible hermitian representations that we realize on the same vector space V, similar as in 3.4. If there existed $\alpha_1>1/2$ such that $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{1/2}\,\chi_{\omega_{E/F},2};\lambda')$ was unitary, then by [Sch14, Lemma 3.6] $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{1/2}\,\chi_{\omega_{E/F},2};\lambda')$ would be unitary for all $\alpha_1>1/2$, in contradiction to [Sch14, Lemma 3.8].

We show that representations in II of Figure 7, page 17, are non-unitary. Let $\alpha_1>1/2$ and $0<\alpha_2<1/2$. Representations $|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1}\times|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2}\rtimes\lambda'$ are irreducible by [Sch14, Th. 4.3] and equal their own Langlands quotient $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2};\lambda')$. Fix $0<\alpha_2<1/2$. $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2};\lambda')$ give a continuous 1-parameter family of irreducible hermitian representations that we realize on the same vector space V. If there existed $\alpha_1>1/2$ such that $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2};\lambda')$ was unitary, then by [Sch14, Lemma 3.6] representations $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2};\lambda')$ would be unitary $\forall\alpha_1>1/2$, in contradiction to [Sch14, Lemma 3.8].

We show that representations in III of Figure 7, page 17, are non-unitary. Let $\alpha_1,\alpha_2>1/2$, $\alpha_2\leq\alpha_1$. The representations $|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1}\times|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2}\rtimes\lambda'$ are irreducible by [Sch14, Th. 4.3] and equal to their own Langlands quotient $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2};\lambda')$. Fix $\alpha_1=\alpha_2=:\alpha$. Let $t\in[1,\infty[$. $\mathrm{Lg}(|\,|^{t\alpha}\,\chi_{\omega_{E/F},1};|\,|^{\alpha}\,\chi_{\omega_{E/F},2};\lambda')$ gives a continuous 1-parameter family of irreducible hermitian representations that we realize on the same vector space V, similar as in 3.4. If there existed $t\in[1,\infty[$ such that $\mathrm{Lg}(|\,|^{t\alpha}\,\chi_{\omega_{E/F},1};|\,|^{\alpha}\,\chi_{\omega_{E/F},2};\lambda')$ was unitary, then by [Sch14, Lemma 3.6] $\mathrm{Lg}(|\,|^{t\alpha}\,\chi_{\omega_{E/F},1};|\,|^{\alpha}\,\chi_{\omega_{E/F},2};\lambda')$ would be unitary $\forall t\in[1,\infty[$, in contradiction to Lemma [Sch14, Lemma 3.8]. Hence $\mathrm{Lg}(|\,|^{\alpha_1}\,\chi_{\omega_{E/F},1};|\,|^{\alpha_2}\,\chi_{\omega_{E/F},2};\lambda')$ is non-unitary for $\alpha_1,\alpha_2>1/2,\alpha_2\leq\alpha_1$.

3. $\chi_{\omega_{E/F},1}\times\chi_{\omega_{E/F},2}\rtimes\lambda'$ is irreducible by [Sch14, Th. 4.1] and unitary. $|\,|^{\alpha}\,\chi_{\omega_{E/F},1}\times\chi_{\omega_{E/F},2}\rtimes\lambda'$ is irreducible for $0<\alpha<1/2$ by [Sch14, Th. 4.5] and equal to its Langlands-quotient $\mathrm{Lg}(|$

$|^{\alpha}\ \chi_{\omega_{E/F},1};\chi_{\omega_{E/F},2}\rtimes\lambda'$). By [Sch14, Lemma 3.6] these Langlands-quotients are unitary. For $\alpha = 1/2$, $|\ |^{1/2}\ \chi_{\omega_{E/F},1}\times\chi_{\omega_{E/F},2}\rtimes\lambda'$ reduces for the first time. By [Mil73] Lg($|\ |^{1/2}\ \chi_{\omega_{E/F},1};\chi_{\omega_{E/F},2}\rtimes\lambda'$) is unitary.

4. For $\alpha > 1/2$, $|\ |^{\alpha}\ \chi_{\omega_{E/F},1}\times\chi_{\omega_{E/F},2}\rtimes\lambda'$ is irreducible by [Sch14, Th. 4.5] and equal to its Langlands-quotient Lg($|\ |^{\alpha}\ \chi_{\omega_{E/F},1};\chi_{\omega_{E/F},2}\rtimes\lambda'$). By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary. (Figure 7, page 17) □




Figure 7

FIGURE 7. Let $\alpha_1,\alpha_2 \geq 0$, let $\chi_{\omega_{E/F},1},\chi_{\omega_{E/F},2} \in X_{\omega_{E/F}}$, such that $\chi_{\omega_{E/F},1} \ncong \chi_{\omega_{E/F},2}$. Figure 7 shows lines and points of reducibility of the representation $|\ |^{\alpha_1}\ \chi_{\omega_{E/F},1}\times\ |\ |^{\alpha_2}\ \chi_{\omega_{E/F},2}\rtimes\lambda'$ and the unitary dual. Lg($|\ |^{\alpha_1}\ \chi_{\omega_{E/F},1};|\ |^{\alpha_2}\ \chi_{\omega_{E/F},2};\lambda'$) is unitary for $0<\alpha_2\leq\alpha_1\leq 1/2$. Otherwise it is non-unitary. Lg($|\ |^{\alpha}\ \chi_{\omega_{E/F},1};\chi_{\omega_{E/F},2}\rtimes\lambda'$) is unitary for $0<\alpha\leq 1/2$. Otherwise it is non-unitary.

$$\mathrm{Lg}(|\ |^{\alpha_1}\ \chi_{1_{F^*},1};|\ |^{\alpha_2}\ \chi_{1_{F^*},2};\lambda'),\ 0<\alpha_2\leq\alpha_1$$

Let $\chi_{1_{F^*},1},\chi_{1_{F^*},2}\in X_{1_{F^*}}$, such that $\chi_{1_{F^*}}\ncong\chi_{2_{F^*}}$.

Theorem 3.14. *Let $\alpha_1,\alpha_2>0$.* Lg($|\ |^{\alpha_1}\ \chi_{1_{F^*},1};|\ |^{\alpha_2}\ \chi_{1_{F^*},2};\lambda'$) *is non-unitary.*

Proof. Representations $|\ |^{\alpha_1}\ \chi_{1_{F^*},1}\times|\ |^{\alpha_2}\ \chi_{1_{F^*},2}\rtimes\lambda'$ are irreducible by [Sch14, Th. 4.3] and equal to their own Langlands-quotient Lg($|\ |^{\alpha_1}\ \chi_{1_{F^*},1};|\ |^{\alpha_2}\ \chi_{1_{F^*},2};\lambda'$). As in 3.4 (2), by [Sch14, Lemmas 3.6 and 3.8], we obtain that Lg($|\ |^{\alpha_1}\ \chi_{1_{F^*},1};|\ |^{\alpha_2}\ \chi_{1_{F^*},2};\lambda'$) is non-unitary. (Figure 8, page 18) □

3.2. **Representations induced from M_1, with cuspidal support in M_0, not fully-induced.**

Irreducible subquotients of $\chi\,\mathrm{St}_{\mathrm{GL}_2}\rtimes\lambda'$

Let χ be a unitary character of E^*.

$\chi\,\mathrm{St}_{\mathrm{GL}_2}\rtimes\lambda'$ is tempered, unitary. Hence all irreducible subquotients of $\chi\,\mathrm{St}_{\mathrm{GL}_2}\rtimes\lambda'$ are tempered and unitary.




Figure 8

FIGURE 8. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{1_{F^*},1}, \chi_{1_{F^*},2} \in X_{1_{F^*}}$ such that $\chi_{1_{F^*},1} \ncong \chi_{1_{F^*},2}$. Figure 8 shows lines and points of reducibility of the representation $|\ |^{\alpha_1}\ \chi_{1_{F^*},1} \times |\ |^{\alpha_2}\ \chi_{1_{F^*},2} \rtimes \lambda'$ and the unitary dual. $\mathrm{Lg}(|\ |^{\alpha_1}\ \chi_{1_{F^*},1}; |\ |^{\alpha_2}\ \chi_{1_{F^*},2}; \lambda')$ is non-unitary for $0 < \alpha_2 \leq \alpha_1$.

Remark 3.15 By [Sch14, Prop. 4.8] $\chi\,\mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ is reducible if and only if $\chi =: \chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$. $\chi_{\omega_{E/F}}\,\mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ has two tempered subquotients.

$$\mathrm{Lg}(|\ |^{\alpha}\ \chi\,\mathrm{St}_{\mathrm{GL}_2}; \lambda'),\ \alpha > 0$$

Theorem 3.16. *Let χ be a unitary character of E^* such that $\chi \notin X_{N_{E/F}(E^*)}$. $\mathrm{Lg}(|\ |^{\alpha}\ \chi\,\mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is non-unitary $\forall \alpha > 0$.*

Proof. If $\chi \notin X_{N_{E/F}(E^*)}$, the representations $\mathrm{Lg}(|\ |^{\alpha}\ \chi\,\mathrm{St}_{\mathrm{GL}_2}; \lambda')$ are not hermitian. □

Theorem 3.17. *1. Let $0 < \alpha \leq 1/2$. $\mathrm{Lg}(|\ |^{\alpha}\ \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is unitary.*
2. Let $\alpha > 1/2$. $\mathrm{Lg}(|\ |^{\alpha}\ \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is non-unitary.

Proof. 1. Let $0 < \alpha < 1/2$. The representations $|\ |^{\alpha}\ \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.7]. They form a continuous one-parameter family of irreducible hermitian representations, that, similar as in 3.4, we realize on the same vector space V. $\mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ is irreducible by [Sch14, Prop. 4.8] and tempered, hence unitary. By [Sch14, Lemma 3.6] the representations $|\ |^{\alpha}\ \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda' = \mathrm{Lg}(|\ |^{\alpha}\ \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ are unitary for $0 < \alpha < 1/2$. By [Sch14, Th. 4.7] $|\ |^{1/2}\ \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ is reducible. By [Mil73] $\mathrm{Lg}(|\ |^{1/2}\ \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is unitary.

2. Let $1/2 < \alpha < 3/2$. The representations $|\ |^{\alpha}\ \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.7] and equal to their Langlands quotient $\mathrm{Lg}(|\ |^{\alpha}\ \mathrm{St}_{\mathrm{GL}_2}; \lambda')$. They form a continuous 1 - parameter family of irreducible hermitian representations, that we realize on the same vector space V. For $\alpha = 3/2$, $\mathrm{Lg}(|\ |^{3/2}\ \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is a subquotient of the representation $|\ |^{2}\ 1 \times |\ |^{1}\ 1 \rtimes \lambda'$ ([Sch14, Th. 5.2]). By results of Casselmann [Cas81], page 915, it is non-unitary. By [Mil73] and [Sch14, Lemma 6.3] $\mathrm{Lg}(|\ |^{\alpha}\ \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is not unitary for $1/2 < \alpha < 3/2$.

The representations $\mathrm{Lg}(|\ |^{\alpha}\ \mathrm{St}_{\mathrm{GL}_2}; \lambda')$, $\alpha > 3/2$, form a continuous 1-paramater family of irreducible hermitian representations. If there existed $\alpha > 3/2$ such that $\mathrm{Lg}(|\ |^{\alpha}\ \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ was unitary, then by [Sch14, Lemma 6.3] $\mathrm{Lg}(|\ |^{\alpha}\ \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ would be unitary for all $\alpha > 3/2$, in contradiction to [Sch14, Lemma 6.8]. (Figure 9, page 19) □




Figure 9

FIGURE 9. Let $\alpha_1, \alpha_2 \geq 0$. Figure 9 shows lines and points of reducibility of the representation $|\ |^{\alpha_1} 1\times |\ |^{\alpha_2} 1 \rtimes \lambda'$. For $0 < \alpha \leq 1/2$, $\mathrm{Lg}(|\ |^{\alpha} \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda')$ is unitary, for $\alpha > 1/2$ it is non-unitary.

Theorem 3.18. *Let* $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$.
Let $\alpha > 1/2$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ *is non-unitary.*

Proof. Let $1/2 < \alpha < 1$. $|\ |^{\alpha} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.7] and equal to its own Langlands quotient $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$. For $\alpha = 1$, by [Sch14, Th. 5.4] $|\ |\ \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ is reducible, by Remark 4.7 in [HT10], $\mathrm{Lg}(|\ |\ \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is non-unitary. By [Mil73] and [Sch14, Lemma 6.3] $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is not unitary for $1/2 < \alpha < 1$.

Let $\alpha > 1$. $|\ |^{\alpha} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.7] and equal to its own Langlands quotient $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$. If there existed $\alpha > 1$ such that $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ was unitary, then by [Sch14, Lemma 6.3] $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ would be unitary $\forall \alpha > 1$, in contradiction to [Sch14, Lemma 6.8]. (Figure 10, page 20) □

Remark 3.19 Let $0 < \alpha < 1/2$. Then $|\ |^{\alpha} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ is irreducible by [Sch14, Th. 4.7] and equal to its own Langlands quotient $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$. If we assume that $|\ |^{1/2} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ is irreducible and equal to $\mathrm{Lg}(|\ |^{1/2} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$, see [Sch14, Remark 5.5], then we can extend the argument that $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is non-unitary to $\alpha > 0$.

Theorem 3.20. *Let* $\chi_{1_{F^*}} \in X_{1_{F^*}}$.
Let $0 < \alpha \leq 1/2$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{1_{F^*}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ *is unitary.*
Let $\alpha > 1/2$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{1_{F^*}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ *is non-unitary.*

Proof. By Proposition [Sch14, Prop. 4.8] $\chi_{1_{F^*}} \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ is irreducible. It is unitary. Let $0 < \alpha < 1/2$. The representations $|\ |^{\alpha} \chi_{1_{F^*}} \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.7] and equal to their own Langlands quotient $\mathrm{Lg}(|\ |^{\alpha} \chi_{1_{F^*}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$. They form a continuous 1-parameter family of irreducible hermitian representations. Similar as in 3.4 we realize them on the same vector space V. By Lemma [Sch14, Lemma 3.6] $\mathrm{Lg}(|\ |^{\alpha} \chi_{1_{F^*}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is unitary for $0 < \alpha < 1/2$. For $\alpha = 1/2$, $|\ |^{1/2} \chi_{1_{F^*}} \mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ reduces for the first time ([Sch14, Th. 4.7]). By [Mil73] $\mathrm{Lg}(|\ |^{1/2} \chi_{1_{F^*}} \mathrm{St}_{\mathrm{GL}_2}; \lambda')$ is unitary.




Figure 10

FIGURE 10. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$. Figure 10 shows lines and points of reducibility of the representation $|\,|^{\alpha_1}\,\chi_{\omega_{E/F}} \times |\,|^{\alpha_2}\,\chi_{\omega_{E/F}} \rtimes \lambda'$. Let $\alpha > 1/2$. $\mathrm{Lg}(|\,|^{\alpha}\,\chi_{\omega_{E/F}}\,\mathrm{St}_{\mathrm{GL}_2};\lambda')$ is non-unitary.




Figure 11

FIGURE 11. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{1_{F^*}} \in X_{1_{F^*}}$. Figure 11 shows lines and points of reducibility of the representation $|\,|^{\alpha_1}\,\chi_{1_{F^*}} \times |\,|^{\alpha_2}\,\chi_{1_{F^*}} \rtimes \lambda'$. For $0 < \alpha \leq 1/2$, $\mathrm{Lg}(|\,|^{\alpha}\,\chi_{1_{F^*}}\,\mathrm{St}_{\mathrm{GL}_2};\lambda')$ is unitary, for $\alpha > 1/2$ it is non-unitary.

Let $\alpha > 1/2$. Representations $|\,|^{\alpha}\,\chi_{1_{F^*}}\,\mathrm{St}_{\mathrm{GL}_2} \rtimes \lambda'$ are irreducible by [Sch14, Th. 4.7] and equal to their own Langlands quotient $\mathrm{Lg}(|\,|^{\alpha}\,\chi_{1_{F^*}}\,\mathrm{St}_{\mathrm{GL}_2};\lambda')$. If there existed $\alpha > 1/2$ such that $\mathrm{Lg}(|\,|^{\alpha}\,\chi_{1_{F^*}}\,\mathrm{St}_{\mathrm{GL}_2};\lambda')$ is unitary, then by [Sch14, Lemma 3.6] all representations $\mathrm{Lg}(|\,|^{\alpha}\,\chi_{1_{F^*}}\,\mathrm{St}_{\mathrm{GL}_2};\lambda')$ would be unitary for $\alpha > 1/2$, in contradiction to [Sch14, Lemma 3.8]. (Figure 11, page 20) □

3.3. **Representations induced from M_2, with cuspidal support in M_0, not fully-induced.**

Irreducible subquotients of $\chi \rtimes \tau$, τ tempered non-cuspidal of $U(3)$, not fully-induced

Recall that $\lambda'(\det)\,\mathrm{St}_{U(3)}$ is the unique square-integrable subquotient of $|\,|\,1 \rtimes \lambda'$ [Key84]. Let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$. Let $\pi_{1,\chi_{\omega_{E/F}}}$ denote the unique square-integrable irreducible subquotient of $|\,|^{1/2}$

$\chi\omega_{E/F} \rtimes \lambda'$. Let $\chi_{1_{F^*}} \in X_{1_{F^*}}$. We have $\chi_{1_{F^*}} \rtimes \lambda' = \sigma_{1,\chi_{1_{F^*}}} \oplus \sigma_{2,\chi_{1_{F^*}}}$, where $\sigma_{1,\chi_{1_{F^*}}}$ and $\sigma_{2,\chi_{1_{F^*}}}$ are irreducible tempered [Key84].

$\lambda'(\det)\,\mathrm{St}_{U(3)}, \pi_{1,\chi_{\omega_{E/F}}}, \sigma_{1,\chi_{1_{F^*}}}$ and $\sigma_{2,\chi_{1_{F^*}}}$ are all non-cuspidal tempered representations of $U(3)$ that are not fully induced [Key84].

Let χ be a unitary character of E^*.

The representations $\chi \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)},\ \chi \rtimes \pi_{1,\chi_{\omega_{E/F}}},\ \chi \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $\chi \rtimes \sigma_{2,\chi_{1_{F^*}}}$ are tempered, hence unitary. Hence all their irreducible subquotients are tempered, hence unitary.

Remark 3.21 By [Sch14, Prop. 4.10] $\chi \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)}$ is reducible if and only if $\chi = 1$ or $\chi \in X_{1_{F^*}}$. By [Sch14, Prop. 4.12] $\chi \rtimes \pi_{1,\chi_{\omega_{E/F}}}$ is reducible if and only if $\chi \in X_{1_{F^*}}$. By [Sch14, Th. 4.1] $\chi \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $\chi \rtimes \sigma_{2,\chi_{1_{F^*}}}$ are reducible if and only if $\chi \in X_{1_{F^*}}$ but $\chi \ncong \chi_{1_{F^*}}$.

Let $\chi_{1_{F^*}} \in X_{1_{F^*}}$. $1 \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)},\ \chi_{1_{F^*}} \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)},\ \chi_{1_{F^*}} \rtimes \pi_{1,\chi_{\omega_{E/F}}},\ \chi \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $\chi \rtimes \sigma_{2,\chi_{1_{F^*}}}$, where $\chi \in X_{1_{F^*}}$ but $\chi \ncong \chi_{1_{F^*}}$ have two tempered subquotients ([Sch14, Propositions 4.10 and 4.12, Th. 4.1]).

$$\mathrm{Lg}(|\ |^\alpha \chi; \tau),\ \alpha > 0,\ \tau \text{ tempered non-cuspidal of } U(3), \text{ not fully-induced}$$

Recall that $\lambda'(\det)\,\mathrm{St}_{U(3)}$ is the unique square-integrable subquotient of $|\ | \rtimes \lambda'$ [Key84]. Let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$. Let $\pi_{1,\chi_{\omega_{E/F}}}$ denote the unique square-integrable irreducible subquotient of $|\ |^{1/2} \chi_{\omega_{E/F}} \rtimes \lambda'$. Let $\chi_{1_{F^*}} \in X_{1_{F^*}}$. We have $\chi_{1_{F^*}} \rtimes \lambda' = \sigma_{1,\chi_{1_{F^*}}} + \sigma_{2,\chi_{1_{F^*}}}$, where $\sigma_{1,\chi_{1_{F^*}}}$ and $\sigma_{2,\chi_{1_{F^*}}}$ are irreducible tempered [Key84].

$\lambda'(\det)\,\mathrm{St}_{U(3)}, \pi_{1,\chi_{\omega_{E/F}}}, \sigma_{1,\chi_{1_{F^*}}}$ and $\sigma_{2,\chi_{1_{F^*}}}$ are all non-cuspidal tempered representations of $U(3)$ that are not fully induced [Key84].

Theorem 3.22. *Let $\chi \notin X_{N_{E/F}}(E^*)$.*

Let $\alpha > 0$. $\mathrm{Lg}(|\ |^\alpha \chi; \lambda'(\det)\,\mathrm{St}_{U(3)}), \mathrm{Lg}(|\ |^\alpha \chi, \pi_{1,\chi_{\omega_{E/F}}}), \mathrm{Lg}(|\ |^\alpha \chi; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^\alpha \chi; \sigma_{2,\chi_{1_{F^*}}})$ are non-unitary.

Proof. The representations $\mathrm{Lg}(|\ |^\alpha \chi; \lambda'(\det)\,\mathrm{St}_{U(3)}), \mathrm{Lg}(|\ |^\alpha \chi, \pi_{1,\chi_{1_{F^*}}}), \mathrm{Lg}(|\ |^\alpha \chi; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^\alpha \chi; \sigma_{2,\chi_{1_{F^*}}})$ are not hermitian, hence not unitary. □

Let $\chi \in X_{N_{E/F}}(E^*) = 1 \cup X_{\omega_{E/F}} \cup X_{1_{F^*}}$.

Let $\chi_{\omega_{E/F}},\ \chi'_{\omega_{E/F}} \in X_{\omega_{E/F}}$, such that $\chi'_{\omega_{E/F}} \neq \chi_{\omega_{E/F}}$, and let $\chi_{1_{F^*}},\ \chi'_{1_{F^*}} \in X_{1_{F^*}}$, such that $\chi'_{1_{F^*}} \neq \chi_{1_{F^*}}$.

Theorem 3.23. *1. Let $\alpha > 1$. $\mathrm{Lg}(|\ |^\alpha 1; \lambda'(\det)\,\mathrm{St}_{U(3)})$ is non-unitary.*

2. Let $0 < \alpha \leq 1$. $\mathrm{Lg}(|\ |^\alpha 1; \pi_{1,\chi_{\omega_{E/F}}}),\ \mathrm{Lg}(|\ |^\alpha 1; \sigma_{1,\chi_{1_{F^*}}}),\ \mathrm{Lg}(|\ |^\alpha 1; \sigma_{2,\chi_{1_{F^*}}}), \mathrm{Lg}(|\ |^\alpha \chi_{1_{F^*}}; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^\alpha \chi_{1_{F^*}}; \sigma_{2,\chi_{1_{F^*}}})$ are unitary.

3. Let $\alpha > 1$. $\mathrm{Lg}(|\ |^\alpha 1; \pi_{1,\chi_{\omega_{E/F}}}),\ \mathrm{Lg}(|\ |^\alpha 1; \sigma_{1,\chi_{1_{F^*}}}),\ \mathrm{Lg}(|\ |^\alpha 1; \sigma_{2,\chi_{1_{F^*}}}), \mathrm{Lg}(|\ |^\alpha \chi_{1_{F^*}}; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^\alpha \chi_{1_{F^*}}; \sigma_{2,\chi_{1_{F^*}}})$ are non-unitary.

4. Let $0 < \alpha \leq 1/2$. $\mathrm{Lg}(|\ |^\alpha \chi_{\omega_{E/F}}; \lambda'(\det)\,\mathrm{St}_{U(3)}), \mathrm{Lg}(|\ |^\alpha \chi_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}}),\ \mathrm{Lg}(|\ |^\alpha \chi'_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}}),\ \mathrm{Lg}(|\ |^\alpha \chi_{\omega_{E/F}}; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^\alpha \chi_{\omega_{E/F}}; \sigma_{2,\chi_{1_{F^*}}})$ are unitary.

5. Let $\alpha > 1/2$. $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \lambda'(\det)\,\mathrm{St}_{U(3)}), \mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi'_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \sigma_{1,\chi_{1_{F^*}}})$ *and* $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \sigma_{2,\chi_{1_{F^*}}})$ *are non-unitary.*

6. Let $\alpha > 0$. $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}; \lambda'(\det)\,\mathrm{St}_{U(3)})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi'_{1_{F^*}}; \sigma_{1,\chi_{1_{F^*}}})$, *and* $\mathrm{Lg}(|\ |^{\alpha}\ \chi'_{1_{F^*}}; \sigma_{2,\chi_{1_{F^*}}})$, *are non-unitary.*

Proof. 1. Let $1 < \alpha < 2$. By [Sch14, Th. 4.9] $|\ |^{\alpha} 1 \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)}$ is irreducible and equal to its Langlands-quotient $\mathrm{Lg}(|\ |^{\alpha} 1; \lambda'(\det)\,\mathrm{St}_{U(3)})$. By [Cas81] the representation $|\ |^{2} 1 \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)}$ is reducible. By the same author $\mathrm{Lg}(|\ |^{2} 1; \lambda'(\det)\,\mathrm{St}_{U(3)})$ is non-unitary. By [Mil73] and [Sch14, Lemma 3.6] $\mathrm{Lg}(|\ |^{\alpha} 1; \lambda'(\det)\,\mathrm{St}_{U(3)})$ is non-unitary for $1 < \alpha < 2$.

Let $\alpha > 2$. $|\ |^{\alpha} 1 \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)}$ is irreducible by [Sch14, Th. 4.9] and equal to its Langlands-quotient $\mathrm{Lg}(|\ |^{\alpha} 1; \lambda'(\det)\,\mathrm{St}_{U(3)})$. By [Sch14, Lemmas 3.6 and 3.8] and $\mathrm{Lg}(|\ |^{\alpha} 1; \lambda'(\det)\,\mathrm{St}_{U(3)})$ is non-unitary for $\alpha > 2$.

Let $0 < \alpha \leq 1$. We have no proof that $\mathrm{Lg}(|\ |^{\alpha} 1; \lambda'(\det)\,\mathrm{St}_{U(3)})$ is non-unitary. (Figure 12, page 23).

2. The representations $1 \rtimes \pi_{1,\chi_{\omega_{E/F}}}$ are irreducible by [Sch14, Prop. 4.12], representations $1 \rtimes \sigma_{1,\chi_{1_{F^*}}}, 1 \rtimes \sigma_{2,\chi_{1_{F^*}}}, \chi_{1_{F^*}} \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $\chi_{1_{F^*}} \rtimes \sigma_{2,\chi_{1_{F^*}}}$ are irreducible by [Sch14, Th. 4.1]. All representations are unitary. For $0 < \alpha < 1$, representations $|\ |^{\alpha} 1 \rtimes \pi_{1,\chi_{\omega_{E/F}}}$ are irreducible by [Sch14, Th. 4.11], representations $|\ |^{\alpha} 1 \rtimes \sigma_{1,\chi_{1_{F^*}}}$, $|\ |^{\alpha} 1 \rtimes \sigma_{2,\chi_{1_{F^*}}}$, $|\ |^{\alpha}\ \chi_{1_{F^*}} \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $|\ |^{\alpha}\ \chi_{1_{F^*}} \rtimes \sigma_{2,\chi_{1_{F^*}}}$ are irreducible by [Sch14, Th. 4.13]. The representations are equal to their own Langlands-quotients $\mathrm{Lg}(|\ |^{\alpha} 1; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{\alpha} 1; \sigma_{1,\chi_{1_{F^*}}})$, $\mathrm{Lg}(|\ |^{\alpha} 1; \sigma_{2,\chi_{1_{F^*}}})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}; \sigma_{2,\chi_{1_{F^*}}})$, respectively. By Lemma [Sch14, Lemma 3.6] these Langlands-quotients are unitary. For $\alpha = 1$, $|\ |\ 1 \rtimes \pi_{1,\chi_{\omega_{E/F}}}$, $|\ |\ 1 \rtimes \sigma_{1,\chi_{1_{F^*}}}$, $|\ |\ 1 \rtimes \sigma_{2,\chi_{1_{F^*}}}$, $|\ |\ \chi_{1_{F^*}} \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $|\ |\ \chi_{1_{F^*}} \rtimes \sigma_{2,\chi_{1_{F^*}}}$ reduce for the first time ([Sch14, Th. 4.11 and 4.13]. By [Mil73] $\mathrm{Lg}(|\ |\ 1; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |\ 1; \sigma_{1,\chi_{1_{F^*}}})$, $\mathrm{Lg}(|\ |\ 1; \sigma_{2,\chi_{1_{F^*}}})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}; \sigma_{2,\chi_{1_{F^*}}})$ are unitary.

3. For $\alpha > 1$, representations $|\ |^{\alpha} 1 \rtimes \pi_{1,\chi_{\omega_{E/F}}}$, $|\ |^{\alpha} 1 \rtimes \sigma_{1,\chi_{1_{F^*}}}$, $|\ |^{\alpha} 1 \rtimes \sigma_{2,\chi_{1_{F^*}}}$, $|\ |^{\alpha}\ \chi_{1_{F^*}} \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $|\ |^{\alpha}\ \chi_{1_{F^*}} \rtimes \sigma_{2,\chi_{1_{F^*}}}$ are irreducible ([Sch14, Th. 4.11 and 4.13] and equal to their Langlands-quotients $\mathrm{Lg}(|\ |^{\alpha} 1; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{\alpha} 1; \sigma_{1,\chi_{1_{F^*}}}), \mathrm{Lg}(|\ |^{\alpha} 1; \sigma_{2,\chi_{1_{F^*}}}), \mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}, \sigma_{2,\chi_{1_{F^*}}})$, respectively. By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary.

4. Representations $\chi_{\omega_{E/F}} \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)}$, $\chi_{\omega_{E/F}} \rtimes \pi_{1,\chi_{\omega_{E/F}}}$, $\chi'_{\omega_{E/F}} \rtimes \pi_{1,\chi_{\omega_{E/F}}}$, $\chi_{\omega_{E/F}} \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $\chi_{\omega_{E/F}} \rtimes \sigma_{2,\chi_{1_{F^*}}}$ are irreducible ([Sch14, Prop. 4.10, 4.12 and Th. 4.1]) and unitary. For $0 < \alpha < 1/2$, representations $|\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)}$, $|\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \pi_{1,\chi_{\omega_{E/F}}}$, $|\ |^{\alpha}\ \chi'_{\omega_{E/F}} \rtimes \pi_{1,\chi_{\omega_{E/F}}}$, $|\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $|\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \sigma_{2,\chi_{1_{F^*}}}$ are irreducible ([Sch14, Th. 4.9, 4.11 and 4.13]) and equal to their Langlands-quotients $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \lambda'(\det)\,\mathrm{St}_{U(3)}), \mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi'_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \sigma_{2,\chi_{1_{F^*}}})$, respectively. By [Sch14, Lemma 3.6] these Langlands-quotients are unitary. For $\alpha = 1/2$, $|\ |^{1/2}\ \chi_{\omega_{E/F}} \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)}, |\ |^{1/2}\ \chi_{\omega_{E/F}} \rtimes \pi_{1,\chi_{\omega_{E/F}}}$, $|\ |^{1/2}\ \chi'_{\omega_{E/F}} \rtimes \pi_{1,\chi_{\omega_{E/F}}}$, $|\ |^{1/2}\ \chi_{\omega_{E/F}} \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $|\ |^{1/2}\ \chi_{\omega_{E/F}} \rtimes \sigma_{2,\chi_{1_{F^*}}}$ reduce for the first time ([Sch14, Th. 4.9, 4.11 and 4.13]. By [Mil73] $\mathrm{Lg}(|\ |^{1/2}\ \chi_{\omega_{E/F}}; \lambda'(\det)\,\mathrm{St}_{U(3)}), \mathrm{Lg}(|\ |^{1/2}\ \chi_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{1/2}\ \chi'_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{1/2}\ \chi_{\omega_{E/F}}; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^{1/2}\ \chi_{\omega_{E/F}}; \sigma_{2,\chi_{1_{F^*}}})$ are unitary.




Figure 12

FIGURE 12. Let $\alpha_1, \alpha_2 \geq 0$. Figure 12 shows lines and points of reducibility of the representation $|\ |^{\alpha_1}\ 1\times\ |\ |^{\alpha_2}\ 1 \rtimes \lambda'$. For $\alpha > 1$, $\mathrm{Lg}(|\ |^{\alpha}\ 1 \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)})$ is non-unitary.

5. For $\alpha > 1/2$, representations $|\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)}, |\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \pi_{1,\chi_{\omega_{E/F}}}$, $|\ |^{\alpha}\ \chi'_{\omega_{E/F}} \rtimes \pi_{1,\chi_{\omega_{E/F}}}$, $|\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $|\ |^{\alpha}\ \chi_{\omega_{E/F}} \rtimes \sigma_{2,\chi_{1_{F^*}}}$ are irreducible ([Sch14, Th. 4.9, 4.11 and 4.13]) and equal to their Langlands-quotients $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \lambda'(\det)\,\mathrm{St}_{U(3)}), \mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi'_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \sigma_{2,\chi_{1_{F^*}}})$, respectively. By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary.

6. Let $\alpha > 0$. The representations $|\ |^{\alpha}\ \chi_{1_{F^*}} \rtimes \lambda'(\det)\,\mathrm{St}_{U(3)}$, $|\ |^{\alpha}\ \chi_{1_{F^*}} \rtimes \pi_{1,\chi_{\omega_{E/F}}}$, $|\ |^{\alpha}\ \chi'_{1_{F^*}} \rtimes \sigma_{1,\chi_{1_{F^*}}}$ and $|\ |^{\alpha}\ \chi'_{1_{F^*}} \rtimes \sigma_{2,\chi_{1_{F^*}}}$ are irreducible ([Sch14, Th. 4.9, 4.11 and 4.13]) and equal to their Langlands-quotients $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}; \lambda'(\det)\,\mathrm{St}_{U(3)})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}; \pi_{1,\chi_{\omega_{E/F}}})$, $\mathrm{Lg}(|\ |^{\alpha}\ \chi'_{1_{F^*}}; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^{\alpha}\ \chi'_{1_{F^*}}; \sigma_{2,\chi_{1_{F^*}}})$. By [Sch14, Lemmas 3.6 and 3.8] these Langlands-quotients are non-unitary. □

3.4. **Representations with cuspidal support in M_1.** Recall that $M_1 \cong \mathrm{GL}(2,E) \times E^1$.

$$\mathrm{Lg}(|\ |^{\alpha} \pi; \lambda'), \alpha > 0,\ \pi \text{ a cuspidal unitary representation of } \mathrm{GL}_2(E)$$

Let π be a cuspidal unitary representation of $\mathrm{GL}_2(E)$. Let $\alpha > 0$. Assume $\exists\ g \in \mathrm{GL}(2,E)$ s. t. $\pi(g) \neq \pi((\overline{g}^t)^{-1})$. Then the induced representations $\pi \rtimes \lambda'$ and $|\ |^{\alpha} \pi \rtimes \lambda'$ are irreducible. $\pi \rtimes \lambda'$ is unitary. $|\ |^{\alpha} \pi \rtimes \lambda'$ is not hermitian $\forall \alpha > 0$, hence not unitary.

Assume $\pi(g) = \pi((\overline{g}^t)^{-1})\ \forall g \in \mathrm{GL}_2(E)$. Then π is obtained by base change lift from $U(2)$ to $\mathrm{GL}(2,E)$, that is by endoscopic liftings from endoscopic data of $U(2)$ to data of $\mathrm{GL}_2(E)$ [Rog90].

Let $\mathrm{G} := U(2)$ and $\widetilde{\mathrm{G}} = \mathrm{Res}_{E/F}\,\mathrm{G} = \mathrm{GL}(2,E)$.

Let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$. Let ${}^L\mathrm{G}$ be the Langlands dual group of G. Recall that σ is defined to be the non-trivial element of $\mathrm{Gal}(E,F)$, i.e. $\sigma(x) = \overline{x}\ \forall x \in E$. Let $\widetilde{\sigma}$ denote the $F-$ automorphism of $\widetilde{\mathrm{G}}$ associated to σ by the $F-$ structure of $\widetilde{\mathrm{G}}$. Let Γ denote the absolute Galois group of E, let




Figure 13

FIGURE 13. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$. Figure 13 shows lines and points of reducibility of the representation $|\ |^{\alpha_1}\ \chi_{\omega_{E/F}} \times |\ |^{\alpha_2}\ \chi_{\omega_{E/F}} \rtimes \lambda'$. Let $\pi_{1,\chi_{\omega_{E/F}}}$ be the unique square-integrable subquotient of $|\ |^{1/2}\ \chi_{\omega_{E/F}} \rtimes \lambda'$. $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \pi_{1,\chi_{\omega_{E/F}}})$ is unitary for $0 < \alpha \leq 1/2$. It is non-unitary for $\alpha > 1/2$.




Figure 14

FIGURE 14. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{1_{F^*}} \in X_{1_{F^*}}$. Figure 14 shows lines and points of reducibility of the representation $|\ |^{\alpha_1}\ \chi_{1_{F^*}} \times |\ |^{\alpha_2}\ \chi_{1_{F^*}} \rtimes \lambda'$. Let $\alpha > 0$. $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}, \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}, \sigma_{2,\chi_{1_{F^*}}})$ are unitary for $0 < \alpha \leq 1$ and non-unitary for $\alpha > 1$.

W_F and W_E denote the Weil groups of F and E, respectively. Let ρ_{G} denote an L-action of Γ on G and let $\rho_{\widetilde{\mathrm{G}}}$ denote an L-action of Γ on $\widetilde{\mathrm{G}}$. One fixes $\omega_\sigma \in W_F \backslash W_E$.

Lemma 3.24 ([Rog90], 4.7)**.** *Up to isomorphisms, the base change problems for $U(2)$ are the endoscopic liftings from endoscopic data $(\mathrm{G}, {}^L\mathrm{G}, 1, \xi)$ and $(\mathrm{G}, {}^L\mathrm{G}, 1, \xi_{\chi_{\omega_{E/F}}})$ for $(\widetilde{\mathrm{G}}, \widetilde{\sigma}, 1)$ to $\widetilde{\mathrm{G}}$. Here*

$$\xi : \quad {}^L\mathrm{G} \ni g \rtimes_{\rho_{\mathrm{G}}} \omega \mapsto (g, g) \rtimes_{\rho_{\widetilde{\mathrm{G}}}} \omega \in {}^L\widetilde{\mathrm{G}}$$




Figure 15

FIGURE 15. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$. Figure 15 shows lines and points of reducibility of the representation $|\ |^{\alpha_1} 1\times |\ |^{\alpha_2} \chi_{\omega_{E/F}} \rtimes \lambda'$. $\mathrm{Lg}(|\ |^{\alpha} 1; \pi_{1,\chi_{\omega_{E/F}}})$ is unitary for $0 < \alpha \leq 1$. It is non-unitary for $\alpha > 1$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{\omega_{E/F}}; \lambda'(\det)\,\mathrm{St}_{U(3)})$ is unitary for $0 < \alpha \leq 1/2$. It is non-unitary for $\alpha > 1/2$.




Figure 16

FIGURE 16. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{1_{F^*}} \in X_{1_{F^*}}$. Figure 16 shows lines and points of reducibility of the representation $|\ |^{\alpha_1} 1\times |\ |^{\alpha_2} \chi_{1_{F^*}} \rtimes \lambda'$. $\mathrm{Lg}(|\ |^{\alpha} 1; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^{\alpha} 1; \sigma_{2,\chi_{1_{F^*}}})$ are unitary for $0 < \alpha \leq 1$ and non-unitary for $\alpha > 1$. $\mathrm{Lg}(|\ |^{\alpha} \chi_{1_{F^*}}; \lambda'(\det)\,\mathrm{St}_{U(3)})$ is non-unitary $\forall \alpha > 0$.

$$\xi_{\chi_{\omega_{E/F}}} : \ {}^L\mathrm{G} \ni g \rtimes_{\rho_{\mathrm{G}}} \omega \mapsto \begin{cases} (g\chi_{\omega_{E/F}}(\omega), g\chi_{\omega_{E/F}}(\omega)) \rtimes_{\rho_{\sim}}{}_{\mathrm{G}} \omega \in {}^L\widetilde{\mathrm{G}} & \text{if } \omega \in W_E \\ (g, -g) \rtimes_{\rho_{\sim}}{}_{\mathrm{G}} \omega_\sigma \in {}^L\mathrm{G} & \text{if } \omega = \omega_\sigma \end{cases}$$

ξ is called standard base change and $\xi_{\chi_{\omega_{E/F}}}$ is called twisted base change ([Kon01]).




Figure 17

FIGURE 17. Let $\alpha_1, \alpha_2 \geq 0$, let $\chi_{\omega_{E/F}} \in X_{\omega_{E/F}}$ and $\chi_{1_{F^*}} \in X_{1_{F^*}}$. Figure 17 shows lines and points of reducibility of the representation $|\ |^{\alpha_1}\ \chi_{\omega_{E/F}} \times |\ |^{\alpha_2}\ \chi_{1_{F^*}} \rtimes \lambda'$. $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \sigma_{1,\chi_{1_{F^*}}})$ and $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{\omega_{E/F}}; \sigma_{2,\chi_{1_{F^*}}})$ are unitary for $0 < \alpha \leq 1/2$ and non-unitary for $\alpha > 1/2$. $\mathrm{Lg}(|\ |^{\alpha}\ \chi_{1_{F^*}}, \pi_{1,\chi_{\omega_{E/F}}})$ is non-unitary $\forall \alpha > 0$.

Let $\Pi_{temp}(\mathrm{G})$ be the set of equivalence classes of irreducible admissible tempered representations of G . Let $\Pi_{temp}(\widetilde{\mathrm{G}})$ be the set of equivalence classes of irreducible admissible tempered representations of $\widetilde{\mathrm{G}}$. Let Π be a tempered L-packet of G, then $\xi(\Pi)$, $\xi_{\chi_{\omega_{E/F}}}(\Pi) \in \Pi_{temp}(\widetilde{\mathrm{G}})$.

As before let π be a cuspidal unitary representation of $\mathrm{GL}(2, E)$. If $\pi(g) = \pi((\overline{g}^t)^{-1})\ \forall g \in \mathrm{GL}(2, E)$, then $\pi = \xi_{\chi_{\omega_{E/F}}}(\Pi)$ or $\pi = \xi(\Pi)$ ([Rog90, 4.2]).

Let π be a cuspidal unitary representation of $\mathrm{GL}_2(E)$.

(1) If $\pi = \xi_{\chi_{\omega_{E/F}}}(\Pi)$, then $\pi \rtimes \lambda'$ is reducible ([Kon01, 4.2]; [Gol94, 6.2]). $\pi \rtimes \lambda' = \tau_1(\pi) + \tau_2(\pi)$, where $\tau_1(\pi)$ and $\tau_2(\pi)$ are irreducible tempered.
$|\ |^{\alpha}\ \pi \rtimes \lambda'$ is irreducible and never unitarisable for $\alpha > 0$ ([Gol94, 6.3]).

(2) If $\pi = \xi(\Pi)$, then $\pi \rtimes \lambda'$ is irreducible ([Kon01, 4.2], ; [Gol94, 6.2]).

By results of Goldberg ([Gol94, 6.3]) one has:
(a) $|\ |^{\alpha}\ \pi \rtimes \lambda'$ is irreducible and unitarisable for $0 < \alpha < 1/2$.
(b) $|\ |^{1/2}\ \pi \rtimes \lambda'$ is reducible. One has $|\ |^{1/2}\ \pi \rtimes \lambda' = \sigma + \mathrm{Lg}(|\ |^{1/2}\ \pi; \lambda')$, where σ is a generic, non-supercuspidal and square-integrable subrepresentation, and $\mathrm{Lg}(|\ |^{1/2}\ \pi; \lambda')$ is unitary.
(c) $|\ |^{\alpha}\ \pi \rtimes \lambda'$ is irreducible and never unitarisable for $\alpha > 1/2$.

We obtain the following

Theorem 3.25. *Let $\pi = \xi_{\chi_{\omega_{E/F}}}(\Pi)$, let $\alpha > 0$. Then*

$\mathrm{Lg}(|\ |^{\alpha}\ \pi; \lambda')$ is non-unitary.

$\pi \rtimes \lambda' = \tau_1(\pi) + \tau_2(\pi)$, where $\tau_1(\pi)$ and $\tau_2(\pi)$ are irreducible tempered.

Let $\pi = \xi(\Pi)$.
Let $0 < \alpha < 1/2$. $\mathrm{Lg}(|\ |^{\alpha}\ \pi, \lambda')$ is unitary.
Let $\alpha = 1/2$. $\mathrm{Lg}(|\ |^{1/2}\ \pi, \lambda')$ is unitary.

Let $\alpha > 1/2$. $\mathrm{Lg}(|\ |^{\alpha}\pi, \lambda')$ is non-unitary.
$\pi \rtimes \lambda'$ is irreducible and unitary.

We have no classification in terms of the Langlands-quotients for the induced representations of $U(5)$ with cuspidal support in $M_2 \cong E^* \times U(3)$. One reason is that we do not have any parametrizatien for the cuspidal representations of $U(3)$.

References

[Aub95] Anne-Marie Aubert, *Dualité dans le groupe de Grothendieck de la catégorie des représentations lisses de longueur finie d'un groupe réductif p-adique*, Trans. Amer. Math. Soc. **347** (1995), no. 6, 2179–2189. MR 1285969 (95i:22025)

[Cas81] W. Casselman, *A new nonunitarity argument for p-adic representations*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **28** (1981), no. 3, 907–928 (1982). MR 656064 (84e:22018)

[Cas95] ———, *Introduction to the theory of admissible representations of p-adic reductive groups*, Draft (1995).

[Gol94] David Goldberg, *Some results on reducibility for unitary groups and local Asai L-functions*, J. Reine Angew. Math. **448** (1994), 65–95. MR 1266747 (95g:22031)

[Gol95] ———, *R-groups and elliptic representations for unitary groups*, Proceedings of the American Mathematical Society **123** (1995), no. 4, pp. 1267–1276 (English).

[Han09] Marcela Hanzer, *The unitarizability of the Aubert dual of strongly positive square integrable representations*, Israel J. Math. **169** (2009), 251–294. MR 2460906 (2009k:22024)

[HT10] Marcela Hanzer and Marko Tadić, *A method of proving non-unitarity of representations of p-adic groups I*, Math. Z. **265** (2010), no. 4, 799–816. MR 2652536 (2011j:22025)

[Key82] Charles David Keys, *On the decomposition of reducible principal series representations of p-adic Chevalley groups*, Pacific J. Math. **101** (1982), no. 2, 351–388. MR 675406 (84d:22032)

[Key84] ———, *Principal series of special unitary groups over local fields*, Compositio Math. **51** (1984), 115–130.

[Kon01] Kazuko Konno, *Induced representations of U(2,2) over a p-adic field*, J. Reine Angew. Math. **540** (2001), 167–204. MR 1868602

[Mil73] Dragan Miličić, *On C^*-algebras with bounded trace*, Glasnik Mat. Ser. III **8(28)** (1973), 7–22. MR 0324429 (48 #2781)

[Mui97] Goran Muić, *The unitary dual of p-adic G_2*, Duke Math. J. **90** (1997), no. 3, 465–493. MR 1480543 (98k:22073)

[Rog90] Jonathan D. Rogawski, *Automorphic representations of unitary groups in three variables*, Annals of Mathematics Studies, vol. 123, Princeton University Press, Princeton, NJ, 1990. MR 1081540 (91k:22037)

[Sch14] Claudia Schoemann, *Irreducible dual of p-adic u(5)*, preprint (2014).

[Tad88] Marko Tadić, *Geometry of dual spaces of reductive groups (non-Archimedean case)*, J. Analyse Math. **51** (1988), 139–181. MR 963153 (90c:22057)

[Tad98] ———, *On reducibility of parabolic induction*, Israel J. Math. **107** (1998), 29–91. MR 1658535 (2001d:22012)

[Tad09] ———, *On reducibility and unitarizability for classical p-adic groups, some general results*, Canad. J. Math. **61** (2009), no. 2, 427–450. MR 2504024 (2010a:22026)